

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



700900

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 48/00, C07K 14/47, C12N 1/15, 1/21, 5/10, 15/09, 15/12, C12P 21/02</b>		<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/31396</b>
			<b>(43) International Publication Date:</b> 23 July 1998 (23.07.98)
<b>(21) International Application Number:</b> PCT/US97/06864		<b>(74) Agent:</b> WILSON, Mary, J.; Nixon & Vanderhye P.C., 8th floor, 1100 North Glebe Road, Arlington, VA 22201-4714 (US).	
<b>(22) International Filing Date:</b> 22 April 1997 (22.04.97)		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
<b>(30) Priority Data:</b> 60/034,960      15 January 1997 (15.01.97)      US		<b>Published</b> <i>With international search report.</i> <i>With an indication in relation to a deposited microorganism furnished under Rule 13bis separately from the description.</i> <i>Date of receipt by the International Bureau:</i> 23 February 1998 (23.02.1998)	
<b>(71) Applicants (for all designated States except US):</b> DUKE UNIVERSITY [US/US]; 230 North Building, Research Drive, Box 90083, Durham, NC 27708-0083 (US). THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 22nd floor, 300 Lakeside Drive, Oakland, CA 94612-3550 (US). CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE CENTRE DE RECHERCHE SUR L'ENDOCRINOLOGIE MOLECULAIRE ET LE DEVELOPPEMENT [FR/FR]; 9, rue Jules Hetzel, F-92190 Muedon (FR).			
<b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> SURWIT, Richard, S. [US/US]; 3804 Sweeten Creek Road, Chapel Hill, NC 27514 (US). COLLINS, Sheila, A. [US/US]; 1516 Woodburn Road, Durham, NC 27705 (US). WARDEN, Craig, H. [US/US]; 728 Waxwing Place, Davis, CA 95616 (US). SELDIN, Michael, F. [US/US]; 3207 Magpie, Davis, CA 95616 (US). RICQUIER, Daniel [FR/FR]; 167, rue Alexis			
<b>(54) Title:</b> RESPIRATION UNCOUPLING PROTEIN			
<b>(57) Abstract</b>  The present invention relates, in general, to a protein linked to cell respiration, thermogenesis, obesity and hyperinsulinemia and, in particular, to a protein designated uncoupling protein-2 (UCP2) and to nucleic acid sequences encoding same. The invention also relates to diagnostic methodologies based, for example, on a determination of levels of UCP2 expression. Further, the invention relates to therapies involving modulating UCP2 expression and/or activity. In addition, the present invention relates to methods of screening compounds for their suitability for use in such therapies.			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**RESPIRATION UNCOUPLING PROTEIN**

The entire content of Provisional Application No. 60/034,960, filed January 15, 1997, is incorporated herein by reference.

**TECHNICAL FIELD**

The present invention relates, in general, to a protein linked to cell respiration, metabolic rate, thermogenesis, obesity and hyperinsulinemia and, in particular, to a protein designated uncoupling protein-2 (UCP2) and to nucleic acid sequences encoding same. The invention also relates to diagnostic methodologies based, for example, on a determination of levels of UCP2 expression. Further, the invention relates to therapies involving modulating UCP2 expression and/or activity. In addition, the present invention relates to methods of screening compounds for their suitability for use in such therapies.

**BACKGROUND**

Obesity is a major health problem in most industrialized countries, including the United States. The seriousness of the pathologies associated with obesity, including type II diabetes and hypertension, has prompted numerous studies directed at determining the molecular basis for obesity. One result of such

studies is the demonstration that genetically inherited characteristics may play a significant role in the development of the disorder.

Recent reports indicate that obesity may be associated with an energy imbalance. The energy balance of an individual is dependent upon both the energy intake (food) and the energy output (corresponding to resting metabolic rate metabolism, physical exercise and heat production). One hypothesis is that decreased resting metabolic rate and/or thermogenesis contributes to the development of obesity. Results of studies relating to uncoupling protein-1 (UCP1) demonstrate that such is the case in rodents (Ricquier et al, FASEB J. 5:2237 (1991); Muzin et al, Biochem. J. 261:721 (1989); Ricquier et al, J. Biol. Chem. 261:13905 (1986); Collins et al, Endocrinol. 138:405 (1997)).

UCP1 creates a pathway that allows dissipation of the electrochemical gradient of protons across the inner mitochondrial membrane in brown adipose tissue, without coupling to any other energy consuming process. This results in generation of heat and dissipation of calories. This process has been implicated in the regulation of body temperature, body composition and glucose metabolism in animals. However, UCP1-containing brown adipose tissue is unlikely to play a role in weight regulation in adult animals and humans living in a thermoneutral

5

10

BRIEF DESCRIPTION OF THE DRAWINGS

25

**SUBSTITUTE SHEET (RULE 26)**

Primers were derived from the mouse UCP2 cDNA to amplify a genomic human fragment corresponding to exons 1 and 2, plus introns 1 and 2 of the human UCP2 gene. This partial genomic fragment was then used to clone the entire human UCP2 gene, hUCP2-g1. The library screened was a human genomic library from Clontech (cat# HL 1067j; lot #45003) made in the EMBL3 SP6/T7 cloning vector with *E. coli* K802 as host strain. The DNA source was human placenta. E. Consensus sequence for the entire human expressed UCP2. The entire expressed sequence (mRNA) is 1612 bp long. The coding sequence extends from bp 345 to 1275 of this sequence. The clones hUCP2-5' and hUCP2-3', from which the consensus sequence was derived, were isolated from the human lung Marathon library from Clontech (Palo Alto, CA; catalog #7408-1). They were isolated by Random Amplification of cDNA Ends (RACE). 5' and 3' RACE were performed on the Marathon library using methods as described by Clontech. The 5' gene specific primer sequence was AGAGAAGGGAAGGAGGGAAG (hUCP2.CDSR3). The 3' gene specific primer sequence was CATCTCCTGGGACGTAGC (hUCP2.CDSF3). PCR conditions were 94°C 30 sec, 64°C 30 sec, 72°C 1 min for 30 cycles on an MJ Research PTC-200, with Clontech KlenTaq Plus DNA polymerase. F. Oligonucleotides that can be used to amplify the UCP2 coding sequence. Names are: hUCP2.CDS (F or R, pair number) where hUCP2 means human UCP2, CDS means coding sequence and F or R

refers to forward or reverse primer. PCR conditions are in an MJ research PTC-200 96-well machine. Clontech KlenTaq plus DNA polymerase were used: 94°C 30 sec, 60°C 30 sec, and 68°C 2 min. Times 30 cycles.

5 Figure 2. Flow cytometry analysis of membrane potential following the expression of UCP2 in *S. cerevisiae*; comparison of wild-type UCP1 and non-active UCP1 mutants. The number of cells (counts, Y-axis) is represented on a logarithmic scale of 1024  
10 channels. The X-axis is a logarithmic scale of fluorescence intensity. A shift of the curve towards the left indicates a decreased mitochondrial potential. The horizontal bar on the left indicates the position of cells treated with mCICCP, a chemical  
15 uncoupler. The expression of UCP2, UCP1 or UCPmut mRNA in recombinant yeast, as well as its absence in wild type yeast, was checked by Northern blots.

Figures 3A-C. A. Tissue distributions of UCP2 in humans and mice. Human UCP2 mRNA was detected on a  
20 human multiple tissue Northern blot containing 2 µg mRNA per lane (Clontech, Palo Alto). This blot was probed with <sup>32</sup>P-labeled human UCP2 insert from IMAGE clone 129216. The <sup>32</sup>P-labeled insert was used to probe the Master blot in ExpressHyb solution (Clontech, Palo  
25 Alto, CA). Expression of UCP1 and UCP2 in mice was examined in various tissues of mice maintained at 23°C. 20 µg RNA was loaded in each lane except the WAT lane which has 4 µg. UCP1 mRNA size is 1.5 kb,

while UCP2 mRNA size is 1.6 kb. The UCP1 probe is the HSU 28480 clone, while UCP2 clone is GenBank accession number U69135 (DR). Abbreviations: H, heart; B, brain; Pl, placenta; Lu, lung; L, liver; M, skeletal muscle; K, kidney; P, pancreas; Wat, white adipose tissue; Bat, brown adipose tissue; and M1 and M2 are thigh and abdominal muscle, respectively.

B. Chromosomal location of Ucp2 on distal mouse chromosome 7. The approximate position in centi-Morgan (cM) distance from the centromere, relationship among other genes and corresponding human chromosomal positions of orthologous genes are shown. The Ucp2 gene segregated in all 114 meiotic events examined with  $\beta$ -arrestin (*Arrb1*), a gene previously mapped to this position on both mouse chromosome 7 and to the long arm of human chromosome 11 (band q13). The other symbols and corresponding names are: *Tyr*, tyrosinase; *D7Was12*, DNA segment Chr 7, Washington 12; *Omp*, olfactory marker protein; *Hbb*, hemoglobin beta chain complex; *tub*, tubby mutation; *Calc*, calcitonin; and *D7Mit40*, DNA segment Chr 7 Mit 40.

C. Regulation of UCP2 mRNA levels in A/J and C57BL/6J by diet. 30  $\mu$ g of epididymal fat pad white adipose tissue RNA from 2 mice from each strain and diet was electrophoresed and a Northern blot prepared. The high-fat diet is the same as described previously (Surwit et al, Diabetes 37:1163 (1988)). The blot was probed with 32P-labeled clone 129216 (human UCP2) and



with cyclophilin. Amounts of UCP2 mRNA were determined with a Molecular Dynamics phosphorimager and were normalized with cyclophilin mRNA levels.

Figure 4. Effect of diet on UCP2 mRNA levels in A/J and B6 mice. Methods are as described in Figure 3C. Epididymal white adipose tissue was prepared from mice on a low fat diet (L), a high fat diet (Surwit et al, Diabetes 37:1163 (1988)) (H), or the high fat diet and the  $\beta 3$  adrenergic agonist CL316,243 (HB) for 25 days (Collins et al, Endocrin. 138:405 (1997)).

Figure 5. Hypothesis about the regulation of resting metabolic rate and thermogenesis by UCP1 and UCP2. In mitochondria, the proton electrochemical gradient ( $\Delta\mu H^+$ ) generated by the respiratory chain allows ATP synthesis by ATP synthase. In brown adipocyte mitochondria, uniquely, UCP1 short-circuits the proton circuit and oxidative energy is dissipated as heat in response to cold exposure. It has also been proposed that cafeteria diets may activate brown adipocyte thermogenesis (Rothwell et al, Nature 281:31 (1979)). In contrast, mitochondria of many tissues express UCP2, an homolog of UCP1, that may also lower the level of coupling of respiration to ADP phosphorylation. This may also lead to increased resting metabolic rate and body temperature and underlies "thermogenesis". The present data show that UCP2 is not induced by cold but is induced by diet.

Figure 6. Increase in UCP2 expression by thiazolidinediones.

Figure 7. Map showing organization of the mouse UCP2 gene.

5

Figures 8A and B. Mouse UCP2 sequences. A. Partial promoter sequence. B. Promoter and coding sequence.

Figure 9. Map showing organization of the human UCP2 gene.

10

Figures 10A-D. Sequences of four regions of the human UCP2 gene.

Figure 11. Functional activity of UCP2 promoter. The following CAT reporter constructs were transfected into HIB-1B cells: SV2 contains the SV40 T-antigen promoter; SV0 is a promoter-less control construct; UCP2(+) and UCP2(-) indicates constructs containing the 246 bp fragment from the mouse UCP2 gene inserted in the "sense" orientation or the "antisense" orientation, respectively. Twenty-four hours following their introduction into HIB-1B, the cells were harvested, a soluble extract of the cells was prepared and the level of CAT activity was measured by the TLC method. The data shown are the mean  $\pm$  sd of two determinations, and the CAT activity is determined as acetylated chloramphenicol product/total, and is expressed as an arbitrary unit. The sample marked as Blank is the unreacted chloramphenicol substrate.

15

20

25

SUMMARY OF THE INVENTION

The present invention relates, in general, to a protein that can effect partial uncoupling of respiration. The protein, designated uncoupling protein-2 (UCP2), is linked to hyperinsulinemia, resting metabolic rate, glucose intolerance, diabetes, obesity, anorexia, cachexia and syndrome X (Reaven, Diabetes 37:1595 (1988); De Franco et al, Diabetes Care 14:173 (1991)). The invention also relates to nucleic acid sequences encoding UCP2 and to diagnostic methodologies based, for example, on a determination of levels of UCP2 expression. Further, the invention relates to therapies involving modulating UCP2 expression and/or activity. In addition, the present invention relates to methods of screening compounds for their suitability for use in such therapies.

Objects and advantages of the present invention will be clear from the description that follows.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to a protein that is involved in energy balance, body weight regulation and thermoregulation. The protein, UCP2, is widely expressed in mammalian tissues, including human tissues, and UCP2 mRNA levels in white fat are elevated in response to fat feeding. The UCP2 gene sequence maps to a chromosomal region linked to obesity and hyperinsulinemia.

5 The identification of UCP2 provides an explanation for a proton leak that has been reported in mitochondria (Brand et al, Biochem. J. 275:81 (1991); Porter et al, Am. J. Physiol. 269:R1213 (1995); Porter et al, Nature 362:628 (1993)). This leak is related to the standard metabolic rate which varies inversely with body mass of various species and with hormonal status (Brand et al, Bio. Chem. J. 275:81 (1991); Porter et al, Nature 362:628 (1993); 10 Rice et al, Obesity Research 4:441 (1996)). Thus, UCP2 proton conductants can be expected to vary with membrane potential, as is the case with UCP1 (Diolez et al, in 8th European Bioenergetic Conference, Valencia, Spain (1994)). This property permits the 15 leak to operate without collapsing membrane potential.

In brown adipose tissue, UCP1 creates a pathway that allows dissipation of the electrochemical gradient of protons across the inner membrane, without coupling to any other energy consuming process 20 (Nedergard et al, in Molecular Mechanisms and Bioenergetics 23:385-420, ed. Ernster, L., Elsevier Science (1992)). The consequences are lower efficiency of oxidative phosphorylation and increased heat production. Previously, this was expected to be 25 a general property of the membrane (involving the phospholipid composition) or a side effect of a known protein such as the ADP/ATP carrier. As shown in Fig. 5, it is now known that UCP2, a thermogenic and

resting metabolic rate protein that is directly regulated by diet, underlies this phenomena. Further, the expression of UCP2 in spleen, bone marrow, macrophages and lymphocytes indicates that this protein also underlies thermogenic responses (e.g. fever) to inflammatory stimuli.

The present invention relates generally to a nucleic acid sequence encoding UCP2, particularly, a mammalian UCP2, more particularly, human UCP2, or portion of that encoding sequence. The invention further relates to the encoded protein, polypeptide or peptide. The term "portion", as used herein, and as applied to nucleic acid sequences, refers to fragments of at least 15 or 30 bases, preferably, at least 50 bases, more preferably, at least 100 bases and, most preferably, at least 150, 300 or 500 bases. As applied to proteins, the term "portion" relates to peptides and polypeptides of at least 5 or 10 amino acids, preferably, at least 17 amino acids, more preferably, at least 33 amino acids and most preferably, at least 50, 100 or 240 amino acids. The invention also relates recombinant molecules comprising the above nucleic acid sequences and to host cells transformed therewith. In addition, the invention relates to methods of making the protein, polypeptide or peptide encoded in the nucleic acid sequence by culturing the transformed host cells under appropriate conditions. Furthermore, the invention

relates to methods of screening compounds for the ability to bind to or alter the activity of or the expression of the UCP2 gene product. In addition, the invention relates to diagnostic and treatment methodologies based on UCP2 and its encoding sequence.

UCP2 Encoding Sequence and Methods of UCP2 Production Using Same, UCP2 and Anti-UCP2 Antibodies

The present invention relates to nucleotide sequences that encode a protein that partially uncouples respiration (as does UCP1) in non brown (as well as brown) adipose tissue, for example, mammalian UCP2, particularly, human UCP2, or portions thereof as defined above (examples of such portions include sequences encoding the 10 N-terminal amino acids, and sequences encoding the mitochondrial carrier protein motifs of Fig. 1A). In particular, the present invention relates to nucleotide sequences that encode the amino acid sequence given in Figure 1A, or portions thereof as defined above (the specific DNA sequence encoding UCP2 given in Figure 1B being only an example). Further, nucleotide sequences to which the invention relates include those encoding substantially the same protein as shown in Figure 1A, for example, inter- and intra-species variations thereof (see Example IX), as well as functional equivalents of the amino acid sequence shown in Figure 1A. The invention further relates to nucleotide

sequences substantially identical to the sequence shown in Figure 1B. A "substantially identical" sequence is one the complement of which hybridizes to the nucleic acid sequence of Figure 1B at 42°C in 50% formamide, 1 X saline/sodium citrate (SSC) containing 0.25% SDS and which remains bound when subjected to washing at 52-55°C with 0.1-0.2 X SSC containing 0.1% SDS (note: 20 X SSC = 3 M sodium chloride/0.3 M sodium citrate). The invention also relates to nucleic acids complementary to those described above.

The present invention also relates to a recombinant molecule comprising a nucleotide sequence as described above and to a host cell transformed therewith. Using standard methodologies, well known in the art, a recombinant molecule comprising a vector and a nucleotide sequence encoding the UCP2 protein, or portion thereof as defined above, can be constructed. Vectors suitable for use in the present invention include plasmid and viral vectors. Plasmid vectors into which a nucleic sequence encoding the UCP2 protein, or portion thereof, can be cloned include any vectors compatible with transformation into a selected host cell. Such vectors include vectors suitable for introduction into yeast and insect cells, generally, mammalian expression vectors suitable for expression in host cells, which vectors can include sequence elements that enhance transcription and/or prolong mRNA halflife in the cell

(e.g.  $\beta$ -globin gene 3' untranslated region)  
specifically, pUC-based *E. coli* vectors, pYeDPUCP2,  
pSelectUCP2, and PECE-UCP2. The nucleotide sequence  
of the invention can be present in the vector operably  
5 linked to regulatory elements, for example, a  
promoter. Suitable promoters include, but are not  
limited to, tissue specific promoters (e.g. leptin  
gene promoter or aP2 gene promoter specific for  
adipose cells, muscle creatine kinase promoter  
10 specific for skeletal muscle and lymphoid cell  
promoters), muscle actin promoter, interleukin  
promoter, CMV, SV40 and MMTV promoters.

As indicated above, the recombinant molecule of  
the invention can be constructed so as to be suitable  
15 for transforming a host cell. Suitable host cells  
include prokaryotic cells, such as bacteria, lower  
eukaryotic cells, such as yeast, and higher eukaryotic  
cells, such as mammalian cells, and insect cells. The  
recombinant molecule of the invention can be  
20 introduced into appropriate host cells by one skilled  
in the art using a variety of known methods.

The present invention further relates to a method  
of producing UCP2, or portions thereof as defined  
above. The method comprises culturing the above-  
25 described transformed host cells under conditions such  
that the encoding sequence is expressed and the  
protein thereby produced.



In addition to the nucleic acids described above, the present invention also relates to UCP2 gene sequences, including introns, exons and flanking regions (e.g. the UCP2 promoter), and to portions thereof suitable for use as probes or primers. The invention also relates to nucleic acid sequences corresponding to the entire expressed UCP2 sequence (e.g. UCP2 mRNA or corresponding cDNA), as well as portions thereof suitable, for example, for use as probes or primers.

A human UCP2 genomic clone, hUCP2-g1, was deposited on January 13, 1997, and was given Accession No. I-1806. A further human genomic clone (designated hUCP2-g2 and a mouse genomic clone (designated MMU2-L2) were deposited at the Collection Nationale de Cultures de Microorganismes, Institut Pasteur, 25, Rue du Docteur Roux, 75724, Paris CEDEX 15, France, under the terms of the Budapest Treaty, on April 16, 1997, and were given Accession Nos. I-1867 and I-1868, respectively. hUCP2-g2 was cloned from the same library used to clone hUCP2-g1. A 500 bp DNA corresponding to the 5' end of hUCP2-g1 was used to screen the genomic library. MMU2-L2 was cloned from a mouse genomic library screened using the mouse UCP2 cDNA. The genomic library was from Strategene (La Jolla, C: 129SVJ Mouse genomic library -- catalog number 946306 -- in the Lamda FIX II vector; the amplification host was XL1-blue MRA(P2). Sequence 2

of hUCP2-g2 is 1161 bp long and goes from position  
-511 to + 650. Therefore, this fragment contains the  
putative human promoter region. A map of the human  
UCP2 gene is shown in Fig. 1C and the sequences of  
several subcloned fragments are given in Fig. 1D.  
(See also Figures 9 and 10.) Two further deposits  
(hUCP2-5' and hUCP2-3') were made at the American Type  
Culture Collection, 12301 Parklawn Drive, Rockville,  
MD 20852, USA, under the terms of the Budapest Treaty,  
on January 10, 1997. These deposits were given  
Accession Nos. 97850 and 97849, respectively. These  
deposits comprise plasmids containing inserts from  
normal human lung cDNA. The plasmids are in the  
Invitrogen pCR2.1 vector. hUCP2-5' includes sequence  
from base pair (bp) 1 up to bp 1375, thus including  
the entire 5' untranslated sequence and the entire  
coding sequence. hUCP2-3' includes sequence from bp  
313 up to bp 1612, thus including the entire coding  
sequence and the entire 3' untranslated region. A  
bacterial artificial chromosome (BAC) clone for human  
genomic UCP2 DNA has been isolated. This clone,  
hUCP2.BAC, was deposited with the American Type  
Culture Collection, 12301 Parklawn Drive, Rockville,  
MD 20852, USA, under the terms of the Budapest Treaty  
on April 18, 1997. The clone was isolated by  
hybridizing hUCP2 probe to the human BAC library  
prepared by Genome Systems. The probe was PCR product  
for human UCP2 produced from human lung cDNA and

Nucleic acid sequence(s) of the invention can be used, in accordance with standard protocols, as probes and primers. Oligonucleotides suitable for amplifying the human UCP2 coding sequence are given in Fig. 1F (pairs 1 and 3 exhibit little non-specific amplification).

The present invention further relates to mammalian UCP2, particularly, human UCP2, substantially free of proteins with which it is normally associated, or portions thereof as defined above. The proteins, polypeptides and peptides of the invention can be produced recombinantly using the nucleic acid sequences as described above, or

chemically using known methods. The protein of the invention can be produced alone or as a fusion product, for example, with a protein such as green fluorescent protein, MaIE protein, glutathione S transferase, glutathione, thrombin, and poly-histidine. Such fusion products can be produced recombinantly. For example, the coding sequence of the invention (e.g. the sequence encoding human UCP2) can be cloned in frame with a sequence encoding another protein (see above examples) the fusion product expressed in an appropriate host cell.

The proteins, polypeptides and peptides of the invention can be used as antigens to generate UCP2 specific antibodies. Methods of antibody generation are well known in the art. Both monoclonal and polyclonal antibodies are included within the scope of the invention, as are binding fragments thereof. One skilled in the art will appreciate that such antibodies can be used to selectively identify and isolate UCP2 and portions thereof. In addition, the antibodies can be used to block activity of UCP2.

## Compound Screens

The present invention also relates to methods of screening compounds for their ability to modulate (e.g. increase or inhibit) the activity or expression of UCP2. More specifically, the present invention relates to methods of testing compounds for their

ability either to increase or to decrease expression or activity of UCP2. The assays are performed *in vitro* or *in vivo*. *In vitro*, cells expressing UCP2 are incubated in the presence and absence of the test compound. By determining the level of UCP2 expression in the presence of the test compound (using, for example, Northern blots, immunoassays (e.g. RIA, Western blots or immunohistochemistry) or PCR), or the level of UCP2 activity in the presence of the test compound, compounds can be identified that suppress or enhance UCP2 expression or activity. Alternatively, constructs comprising the UCP2 promoter operably linked to a reporter gene (e.g. luciferase, chloramphenicol acetyl transferase, LacZ, green fluorescent protein, etc.) can be introduced into host cells and the effect of the test compounds on expression of the reporter gene detected.

Cells suitable for use in the foregoing assays include, but are not limited to, lymphoblasts, myocytes, adipocytes and hepatic cells, more specifically, C2C12 cells, 3T3 cells of adipocyte lineage, HIB-1B cells, rodent hepatoma cells, HepG2 cells, and B7 cells. (See Example V.)

Compounds that suppress or enhance UCP2 expression can also be identified using *in vivo* screens. In these assays, the test compound is administered (e.g. IV, IP, IM, orally, or otherwise), to the animal, for example, at a variety of dose

Compounds that suppress or enhance UCP2 activity can be identified by contacting UCP2 with the test compound under conditions such that the compound can interact with (e.g. bind to) the protein. A system such as the yeast expression system described in

5

10

15

20

25

used to induce hypothermia, for example, when  
advantageous in surgical settings, including  
transplantation. Such agents can also be used to  
block hyperthermia, for example, during thyroid storm.  
5 Compounds that enhance UCP2 expression or stimulate  
its activity can also be used to treat hypothermia.  
Given the similarity in the patterns of leptin and  
UCP2 expression, agents that modulate UCP2 can be  
expected to modulate leptin expression. However,  
10 leptin has been shown not to influence UCP2 expression  
*in vitro* or *in vivo*.

#### Pharmaceutical Compositions

The present invention also relates to  
15 pharmaceutical compositions comprising, as active  
agent, the proteins, peptides, nucleic acids or  
antibodies of the invention. The invention also  
relates to compositions comprising, as active agent,  
compounds selected using the above-described screening  
20 protocols. Such compositions include the active agent  
in combination with a pharmaceutically acceptable  
carrier. In the case of naked DNA, the "carrier" can  
be gold particles. In the case of compositions to be  
applied topically, the composition can take the form  
25 of a gel, cream, ointment or lotion. The amount of  
active agent in the composition can vary with the  
agent, the patient and the effect sought. Likewise,



the dosing regimen can vary depending on the composition and the disease/disorder to be treated.

Detection/Diagnosis:

5           The present invention further relates to methods of identifying individuals at increased risk for developing certain diseases/disorders, including hyperinsulinemia, glucose intolerance, type II diabetes, obesity, syndrome X, immunological  
10           dysfunction and body temperature dysfunction. One such method comprises: (a) obtaining from a mammal (e.g. a preobese human) a biological sample, (b) detecting the presence in the sample of a UCP2 gene product and (c) comparing the amount of the gene  
15           product present in the sample with that in a control sample. Advantageously, the biological sample is taken after the consumption of a high fat meal. In accordance with this method, the presence in the sample of altered (e.g. diminished) levels of UCP2  
20           gene product indicates that the subject is predisposed to the above-indicated diseases/disorders.

          Biological samples suitable for use in this method include biological fluids such as blood. Tissue samples (e.g. biopsies) can also be used in the  
25           method of the invention, including samples derived from muscle or fat. Cell cultures or cell extracts derived, for example, from tissue biopsies can also be used.

The detection step of the present method can be effected using standard protocols for protein/mRNA detection. Examples of suitable protocols include Northern blot analysis, immunoassays (e.g. RIA, Western blots, immunohistochemical analyses), and PCR.

The present invention also relates to methods of identifying individuals having elevated or reduced levels of UCP2, which individuals are likely to benefit from therapies designed to suppress or enhance UCP2 expression, respectively. As indicated above, a biological sample from a preobese subject can be screened for the presence of diminished levels of UCP2 gene product, particularly in response to high fat intake, the presence of depressed levels of the gene product, relative to a normal population (standard), being indicative of predisposition to obesity, type II diabetes or syndrome X. Such individuals would be candidates for anti-obesity therapy (e.g. treatment with appetite suppressants). The identification of elevated levels of UCP2 in a wasting patient (e.g. a cancer, AIDS or anorexia patient) would be indicative of an individual that would benefit from treatment with agents that suppress UCP2 expression or activity. The identification of low levels of UCP2 in a hypothermic patient or obese patient would be indicative of an individual that would benefit from agents that induce UCP2 expression or activity.

5 The present invention also relates to a kit that can be used in the detection of UCP2 expression products. The kit can comprise a compound that specifically binds UCP2 (e.g. binding proteins (e.g. antibodies or binding fragments thereof (e.g. F(ab')<sub>2</sub> fragments)) or UCP2 mRNA (e.g. a complementary probe or primer), for example, disposed within a container means. The kit can further comprise ancillary reagents, including buffers and the like.

10 The diagnostic methodologies described herein are applicable to both humans and non-human mammals.

#### Therapy:

15 The present invention relates to methods of treating diseases/disorders such as hyperinsulinemia, glucose intolerance, diabetes, obesity, syndrome X, cancer and hypothermia by increasing UCP2 activity and/or expression. The invention also relates to methods of treating inflammation, anorexia and wasting (cachexia) (e.g. associated with cancer or AIDS), of  
20 reducing fever and blocking hyperthermia (e.g. thyroid storm) and to methods of inducing hypothermia (eg when advantageous for surgery and transplant), by decreasing UCP2 activity and or expression. These  
25 methodologies can be effected using compounds selected using screening protocols such as those described above and/or by using the gene therapy and antisense approaches described below.

Gene therapy can be used to effect targeted expression of UCP2, for example, in fat tissue and muscle to reduce fat depots or in cancer cells to cause thermodestruction or metabolic collapse/death of the cells. The UCP2 coding sequence can be cloned into an appropriate expression vector and targeted to a particular cell type(s) to achieve efficient, high level expression. Introduction of the UCP2 coding sequence into target cells can be achieved, for example, using particle mediated DNA delivery (William et al, Proc. Natl. Acad. Sci. USA 88:2726 (1991); Haynes, Advances in Drug Delivery Reviews 21:3 (1996)), direct injection of naked DNA (Levy et al, Gene Therapy 3:21 (1996)), liposomal DNA transport (Felgner, Human Gene Therapy 1:1791 (1996)), or viral vector mediated transport (Smith et al, Gene Therapy 3:190 (1996)). Tissue specific effects can be achieved, for example, in the case of virus mediated transport by using viral vectors that are tissue specific, or by the use of promoters that are tissue specific (e.g. leptin and  $\alpha$ 2 promoters can be used to achieve expression in white adipose tissue and the myosin light chain kinase promoter can be used to achieve expression in skeletal muscle) (see also Warden et al, In Regulation of Body Weight: biological and behavioral mechanisms, C. Bouchard and G.A. Bray, eds. (West Sussex; John Wiley & Sons Ltd.), pp. 285-305). Combinatorial approaches can also be used to

ensure that the UCP2 coding sequence is activated in the target tissue (Butt et al, Gene Exp. 4:319 (1995); Hart, Semin. Oncol. 23:1521 (1996)).

5 Antisense oligonucleotides complementary to UCP2 mRNA can be used to selectively diminish or ablate the expression of the protein, for example, at sites of inflammation. More specifically, antisense constructs or antisense oligonucleotides can be used to inhibit the production of UCP2 in high expressing cells  
10 (spleen, thymus, leucocytes, bone marrow and stomach). Antisense mRNA can be produced by transfecting into target cells an expression vector with the UCP2 gene sequence, or portion thereof, oriented in an antisense direction relative to the direction of transcription.  
15 Appropriate vectors include viral vectors, including retroviral, adenoviral, and adeno-associated viral vectors, as well as nonviral vectors. Tissue specific promoters can be used (e.g. leptin gene promoter or aP2 gene promoter specific for adipose cells, muscle  
20 creatine kinase promoter specific for skeletal muscle and lymphoid cell promoters). Alternatively, antisense oligonucleotides can be introduced directly into target cells to achieve the same goal. (See also other delivery methodologies described above in  
25 connection with gene therapy.) Oligonucleotides can be selected/designed to achieve a high level of specificity. (See also Matteucci et al, Nature 384:20 (1996)).

5 It has been recently demonstrated that increased O<sub>2</sub> consumption associated with cachexia of malignancy can be attenuated by indomethacin, a cyclooxygenase inhibitor. This is thought to be due to inhibition of prostaglandin production (Roe et al, Metabolism 46:359 (1997)). Thus, agents that block UCP2 expression and/or activity can be expected to be useful in the treatment of cachexia.

10 The therapeutic methodologies described herein are applicable to both human and non-human mammals (including cats and dogs).

Certain aspects of the present invention are described in greater detail in the non-limiting Examples that follow.

#### 15 EXAMPLES

The following experimental methodologies relate particularly to Examples I and III-VII that follow.

20 Yeast transfections and functional analysis using flow cytometry. The diploid yeast (*Saccharomyces cerevisiae*) strain W303 (a/ $\alpha$ ; ade2-10; his 3-11-15; leu2-2, 112; ura3-1; can1-100; try- $\Delta$ 1) was used for expression of UCP2. The pYedP-UCP2 or UCPmut  
25 expression vectors were introduced into yeast by electroporation and transformants were selected for uracil auxotrophy as described by Bouillaud et al (EMBO J. 13:1990 (1994)). Expression of the UCP2,

UCP1 or UCPmut under the control of the gal-cyc promoter was induced by galactose in the absence of glucose. Growth of yeast for flow cytometry analysis and measurement of mitochondrial membrane potential using the DiOC6(3) fluorescent probe (3,3'-dihexyloxacarbocyanine iodine, Molecular Probes, Eugene, OR) are described by Bouillaud et al (EMBO J. 13:1990 (1994)).

Isolation of UCP2 cDNA clones. Human and mouse UCP2 IMAGE consortium clones (Washington University, St. Louis - Merck) identified in the EST database (<http://www.ncbi.nlm.nih.gov>) were purchased (Research Genetics, Huntsville, Alabama). Human UCP2 sequence was derived from IMAGE clones 129216, 158373, 163246 and 236034. GenBank accession for the human UCP2 coding sequence is U76367. Mouse UCP2 was also isolated independently (DR) by screening a mouse muscle cDNA library with rat UCP1 (GenBank accession U69135). More than 20 mouse and human UCP2 clones are currently in the EST database, while only one UCP1 clone (IMAGE 529189) is represented in the EST database.

Mouse chromosomal mapping. The chromosomal position was determined by linkage mapping of restriction fragment length polymorphisms (RFLPs) in an interspecific cross as previously described (Watson et al, Mamm. Genome 2:158 (1992)). Hybridization of human UCP2 probe (892 bp insert of IMAGE clone 129216)

defined EcoRI RFLPs that segregated in the strains used (C3H, 5.5 kb; *M. spretus*, 9.0 kb). Gene order was determined by analyzing all haplotypes and minimizing crossover frequencies. The reference loci in this interspecific cross that includes 2000 chromosomal markers are indicated with citations in an online database (<http://www.informatics.jax.org/crossdata.html>: select "seldin cross" and chromosome number 7).

#### EXAMPLE I

##### Sequence and Uncoupling Activity of UCP2

The amino acid sequence of human UCP2 is 59% identical to human UCP1 (UCP2 sequence shown in Figure 1A). The predicted coding sequence produces a protein of 309 amino acids with a molecular weight of 33,218 Daltons and an isoelectric point of 10.0. The amino acid sequence of mouse UCP2 (GenBank U69135) is 95% identical to human UCP2. Several protein motifs are conserved between UCP1 and UCP2. Both exhibit three mitochondrial carrier protein motifs, consistent with roles as ion transporters of the inner membrane, while the amino acids essential to ATP binding are also conserved.

To test the hypothesis that UCP2 is an uncoupling protein, yeast were transfected with UCP2 in an expression vector as previously reported for UCP1 (Bouillaud et al, EMBO J. 13:1990 (1994)). Rates of



growth in liquid medium of transformed yeast were measured in the presence of galactose, which induces expression. Instantaneous generation times were compared after induction of vector, UCP2, control (empty) and UCP1 expression vectors. The values for UCP2 were  $260 \pm 29$  minutes ( $\pm$ SD,  $n=5$ ), whereas the mean value for the strain transfected control vector (pYeDP vector) is  $188 \pm 13$  minutes ( $n=5$ ,  $p < 0.01$  versus UCP2) and the mean value for UCP1 is  $226 \pm 10$  minutes ( $n=10$ ,  $p < 0.01$  versus UCP2). These results indicate that UCP2 slows the growth of yeast more than does UCP1 (Bouillaud et al, EMBO J. 13:1990 (1994)).

The *in vivo* decrease of membrane potential associated with uncoupling of respiration was analyzed by flow cytometry of yeast labeled with the potential sensitive probe DiOC6(3). Using this procedure, it was previously demonstrated that UCP1 decreases the mitochondrial potential of transformed yeast (Bouillaud et al, EMBO J. 13:1990 (1994)).

Mitochondrial potential is strongly decreased in strains containing either the UCP2 or UCP1 expression vectors, in contrast to strains containing empty or UCPmut (A36P/K150L mutations that inhibit UCP1 activity) expression vectors (Figure 2). In the case of UCP2, 2 peaks were observed, one equivalent to cells expressing UCP1 and a second peak corresponding to a very low mitochondrial potential. Mean fluorescence intensities were: empty vector, 65.5;

UCPmut, 43.0; UCP1, 22.5; UCP2, 15.5 and UCP2 + mCICCP, 0.4 (Figure 2). Very similar results were obtained in two additional experiments. Results of a representative experiment are shown in Figure 2. These data show that UCP2 influences mitochondrial activity and can partially uncouple respiration.

#### EXAMPLE II

##### Organization of Mouse and Human UCP2 Genes

A map showing the organization of the mouse UCP2 gene present in MMU2-L2. The mouse DNA inserted in lambda phage is 13.9 kb long and contains all the 8 exons and introns, and 5.3 kb of DNA upstream of the putative transcriptional start site (+1 site). (See Fig. 7) The DNA has been sequenced from -934 to +8600 bp (Fig. 8B). The region -246/+1 of the mouse UCP2 gene was placed in front of a CAT reporter gene. This CAT construct exhibited a promoter activity when transfected in cultured adipocytes (see Example X). A fragment of 1.8 kb located in the 5' end of the gene (Fig. 8A) has also been sequenced.

A map showing the organization of the human UCP2 gene present in hUCP-g2. The human DNA inserted in lambda EMBL3 phage is 14 kb long (see Fig. 9). It contains all the 8 exons and introns, and a minimum of 3 kb of DNA upstream of the putative +1 site. Four regions, referred to as sequences 1, 2, 3 or 4 in

Figures 10A-D., have been sequenced. Sequence 1 corresponds to 640 bp of DNA forming the 5' extremity of the human DNA. Sequence 2 corresponds to a 1161 bp DNA from positions bp -511 to +650. This fragment contains the putative proximal human UCP2 promoter.

Human and mouse UCP2 genes have a very similar organization, exons 1 and 2 being noncoding exons, whereas exons 3-6 are coding exons.

### EXAMPLE III

#### **Expression of UCP2 in Human and Mouse Tissues**

To determine the possible sites of physiologic function of UCP2, Northern blot analyses were performed. UCP2 tissue distribution is markedly different from that of UCP1 (Figure 3A). Probing of a multiple tissue northern blot from pooled adult tissues reveals UCP2 mRNA of 1.6 kb size present in skeletal muscle, lung, heart, placenta and kidney. In mice, UCP2 is expressed in brown adipose tissue (BAT) as well as white adipose tissue (WAT), and at high levels in heart and kidney. Probing of a Human mRNA Master Blot (Clontech, Palo Alto, CA) with UCP2 reveals that it is expressed at high levels in skeletal muscle, spleen, thymus, leukocytes, bone marrow and stomach. Low levels of UCP2 mRNA expression were observed in liver or brain.

EXAMPLE IV**Mapping of UCP2 in Mice and Humans**

Additional evidence for the role of UCP2 in obesity and diabetes was sought by determining the chromosomal position of UCP2. Whole genome mapping of mouse genetic crosses has identified at least 10 chromosomal loci influencing spontaneous and diet-induced obesity (Horvat et al, Genetics 139:1737 (1995); Collins et al, Mamm. Genome 4:454 (1993); Warden et al, J. Clin. Invest. 95:1545 (1995); West et al, J. Clin. Invest. 94:1410 (1994); Taylor et al Genomics 34:389 (1996); Seldon et al, J. Clin. Invest. 94:269 (1994)). Identification of the specific genes underlying these loci remains a challenge. A positional candidate approach, in which genes with suspected roles in obesity and insulin resistance are shown to have chromosomal loci co-incident with known obesity and diabetes loci, has been used to show that UCP2 is such a novel candidate gene. UCP2 (*Ucp2*) was mapped to mouse chromosome 7, tightly linked to the *tubby* mutation (Figure 3B). Two independent sequence tagged sites derived from human UCP2 clones (Expressed sequence tags, EST 143230 and 226515) have been mapped to the homologous region of the long arm of human chromosome 11, between D11S916 and D11S911 (WI-1672 and WI-13873, Whitehead Institute Center for Genome Research radiation hybrid panel, <http://www.ncbi.nlm.nih.gov/SCIENCE96/>), consistent

### EXAMPLE V

The chromosomal mapping of *UCP2* is co-incident with quantitative trait loci (QTLs) for obesity from at least three independent mouse models, one congenic strain, and human insulin dependent diabetes locus-4 (IDDM4) (Warden et al, J. Clin. Invest. 95:1545 (1995); Taylor et al, Genomics 34:389 (1996); Seldin et al, J. Clin. Invest. 94:269 (1994); Hashimoto et al, Nature 371:61 (1994)). Diet-induced obesity and diabetes have been demonstrated in the C57BL/6J (B6) mouse, while the A/J strain is resistant to the high-fat diet (Surwit et al, Diabetes 37:1163 (1988)). This syndrome is not due to inactivity (Brownlow et al, Physiology & Behavior 60:37 (1996)) or to hyperphagia (Surwit et al, Metabolism 44:545 (1995)) as B6 mice show higher feed efficiency than A/J mice (weight gained/calories consumed), when fed a high fat diet. Whole genome mapping has shown that a QTL for plasma leptin, insulin, and glucose is located on mouse chromosome 7, peaking near the *tubby* locus (Surwit et al, Diabetes 45:78A (1996)). Sequence analysis and mRNA expression levels suggest that *tubby*

is not the underlying gene for this QTL. When Northern blots of white adipose tissue from A/J and B6 mice were hybridized with the UCP2 probe, mRNA levels were found to be higher in the obesity-resistant A/J than the obesity prone B6 (Figure 4). In addition, UCP2 mRNA levels were significantly increased by the high-fat diet in A/J mice at early time points in the diet (days 7, 18 and 25), but only modestly in B6 animals (Figure 3C). Core body temperature measured by a rectal temperature probe is also increased in A/J and B6 mice consuming the high fat diet, but to a greater extent (2X) in A/J than in B6 mice. These data are highly significant ( $n=10$  for each group,  $p<0.01$ ). However, after progression to severe obesity in the B6 mouse on a high fat diet for 18 weeks, levels of UCP2 were dramatically increased in both strains, but in contrast to early time points (7, 18, 25 days) levels of UCP2 were greater in the diet-induced obese B6 mice relative to the leaner A/J strain. Additionally, core body temperature increased more in A/J mice than in B/6 mice following the introduction of a high fat diet ( $p<0.05$ ). Similar over expression of UCP2 is observed in adult, genetically obese ob/ob and db/db mice.

Regulation of UCP2 has been examined for factors that would regulate UCP1. In 11-week-old fed male Swiss mice, UCP2 mRNA was detected in BAT, WAT, thigh muscle and liver. Exposure at 4°C or treatment with

the  $\beta 3$  adrenergic agonist CL316,243 for 10 days did not influence UCP2 mRNA levels in these tissues. Finally, administration of leptin (2  $\mu$ g, 2x/day for 4 days) failed to increase UCP2 expression. Leptin has been shown to increase sympathetic outflow to brown fat, thus potentially increasing UCP2 expression. Thus, UCP2 is regulated very differently than UCP1, which is induced by cold and  $\beta 3$  agonists. However, these data do not rule out the possibility that the sympathetic nervous system regulates UCP2.

#### EXAMPLE VI

##### **Increase in UCP2 Expression by Thiazolidinediones**

Thiazolidinediones are known insulin-sensitizing agents that lower plasma glucose levels, and long chain fatty acids have been shown to be ligands for the PPAR (peroxisome proliferator activated receptor) family of receptors. Accordingly, a study was undertaken to determine whether agents that stimulate PPAR $\gamma$  could increase expression of UCP2 in a model adipocyte cell line, HIB-1B.

HIB 1B cells were grown for 7 days in DMEM + 10% charcoal-stripped serum (*Media*) or with the addition of the thiazolidine dione BRL 49653 (1 $\mu$ M) and the RXR $\alpha$  ligand LGD-1069 (0.1 $\mu$ M) (TZD/RXR). The results are the average of 2 independent experiments of 4 samples each. \*, significantly different from *Media* samples,  $p < 0.001$ . (See Fig. 6).

EXAMPLE VII**Respiratory Rates**

Respiratory rates have been measured in mitochondria isolated from yeast containing the empty pYeDP vector and the pUCP2 vector (Table 1). Upon addition of the FCCP uncoupler, the respiratory rate was increased by a factor of 7.1 in control mitochondria, while it was only increased by a factor of 4.3 in pUCP2 mitochondria (Table 1). This result clearly demonstrates that UCP2 has uncoupling activity. Addition of ADP stimulates respiration more in control mitochondria, again showing the UCP2 mitochondria are more uncoupled.

Table 1

	Basal rate	Uncoupled		ADP addition	
		rate	fold	rate	fold
pYeDP	14	100	7.1	33	2.3
pUCP2	23	100	4.3	42	1.8

EXAMPLE VIII**Role of UCP2 in a mouse model of spontaneous obesity**

The *Ucp2* gene is co-incident with a QTL for spontaneous multifactorial obesity in BSB mice. BSB mice are derived from a backcross of (C57BL/6J x *Mus spretus*) F1s x C57BL/6J. A locus with significant QTLs for body fat percent, hepatic lipase activity and



plasma cholesterol is located on distal mouse  
chromosome 7 surrounding the *tubby* locus. mRNA  
extracted from the spleens of 16 BSB mice was probed  
on northern blots for UCP2 and actin. Levels of mRNA  
were quantitated by phosphorimager and UCP2 was  
normalized with actin. A negative correlation of  
UCP2/actin mRNA levels with body fat ( $p=0.016$ ) and HDL  
cholesterol ( $p=0.023$ ) was observed. The hypothesis  
that body temperature is correlated with body fat in  
BSB mice was also investigated. Core body  
temperatures were determined by rectal thermometers  
every week after weaning of 45 BSB mice. These BSB  
mice demonstrated a negative correlation of body  
temperature with percent body fat at sacrifice.  
P-values for the correlation of were  $p=0.018$  at 6  
weeks old ( $r^2 = 0.12$ ) and  $p=0.09$  at week 8 ( $r^2=0.10$ ).  
These results indicate that UCP2 influences body fat  
percent by altering body temperature in a model of  
spontaneous obesity exhibited by mice maintained on a  
chow diet. (See Fisler et al, Obesity Res. 1:271  
(1993); Warden et al, J. Clin. Invest. 92:773 (1993);  
Warden et al, J. Clin. Invest. 95:1545 (1995).)

#### EXAMPLE IX

##### Common Sequence Polymorphisms of Human UCP2

Human mRNA was prepared from buffy coats (white  
blood cells) from 8 Hispanics who either have Type II  
diabetes or are the children of Type II diabetics.

5 The complete human UCP2 coding sequence was amplified  
by RT-PCR of mRNA using primer pair 3 (hUCP2.cds3) as  
described in the Brief Description of Fig. 1E. The  
products were cloned into the Invitrogen pCR2.1 vector  
as described by the manufacturer. Sequencing was  
performed with 8 primers on each of the clones.  
Sequences were compared with the hUCP2 sequence  
submitted to Genbank (Accession No. U76367).  
10 5 polymorphisms were observed in at least 2 of the 8  
people sequenced. Each of these confirmed  
polymorphisms was also observed with all overlapping  
primers. 5 additional polymorphisms were observed  
with only one person or have conflicting data from  
different primer pairs. (See Table 2) Restriction  
15 enzymes have been identified that would differentially  
digest each of the alleles.

TABLE 2

Polymorphism	1	2	3	4	5
base number in terms of entire dna sequence	445	502	508	869	913
amino acid number in terms of open reading frame seq.	34	53	55	104	190
hUCP2.genbank	L	A	A	K	N
Confirmed polymorphisms, observed at least twice.	L	A	V	K	S
Polymorphisms observed only once.	P	V	V	K	S
Short name			A55V		N190S

Polymorphism	6	7	8	9	10
base number in terms of entire dna sequence	1052	1068	1142	1224	1262
amino acid number in terms of open reading frame seq.	236	242	266	294	306
hUCP2.genbank	D	Y	P	L	E
Confirmed polymorphisms, observed at least twice.	D	Y	P	M	E
Polymorphisms observed only once.	E	N	P	M	E
short name				L294M	

*Restriction enzymes of HUCP2.genbank for polymorphism analysis:*

Polymorphism 1: Frequency of base change T to C at 445 of entire expressed is 77% T and 33% C. Amino acid change occurs in polymorphisms observed only once from L to P.

site= t/ccgga, possible enzymes= AccIII, BseAI, BseAI, BsiMI, Bsp13I, BspEI, Kpn2I, MroI, There are no other areas where this enzyme would cut.

Polymorphism 2: Frequency of base change C to T at 502 of entire expressed sequence is 22% C, 88% T; base change T to A at 503 of entire expressed sequence is 64% T, 36% A. Amino acid change occurs in edited all sequence from A to V. Restriction enzyme cleavage site= /gtnac, possible enzyme is MaeIII. Other areas that this enzyme cuts are: 119, 583, 865.

Polymorphism 3: Frequency of base change C to T at 508 of entire expressed sequence is 57% C and 43% T as shown in 2 primers. Amino acid change occurs in edited and edited all sequences from A to V. Restriction enzyme cleavage site=gcn/ngc, possible enzyme is Cac8I. Other areas that this enzyme cuts are: 58, 70, 153, 250, 254, 336, 340, 364, 670, 756, 901.

Polymorphism 4: Frequency of base change C to T at 869 of entire expressed sequence is 73% C and 27% T, shown in 2 primers. Seventy-five percent of examined individuals are T. No amino acid change occurs at this possible polymorphism site. Restriction enzyme cleavage site= cctc, possible enzyme is MnlI. Other areas that this enzyme cuts are: 97, 231, 278, 357, 546, 594, 609, 624, 639, 695, 796, 824, 891.

Polymorphism 5: Frequency of base change A to G at 913 of entire expressed sequence is 80% A and 20% G as

shown in two primers. Twenty-five percent of examined individuals are G. Amino acid change occurs in edited and edited all sequences from N to S. Restriction enzyme cleavage site=cmg/ckg where m=a,c and k=g,t; possible enzymes= MspAII, NspBII or site= cag/ctg with enzyme PvuII. Other areas that cmg/ckg cuts are:cat/ctg: 65, 877; ccg/cgg: 177.

Polymorphism 6: Base change C to A at 1052 of entire expressed sequence is included in edited all sequence because 4 out of 8 exhibit A in primer -21M13 and C in primer 241.1. Amino acid change occurs in edited all sequence from D to E. Restriction enzyme cleavage site: gacn/nngtc, possible enzymes are AspI, AtsI, Tth111I. There are no other areas where this site occurs.

Polymorphism 7: Base change T to A at 1068 of entire expressed sequence is included in edited all sequence because 5 out of 8 exhibit A in primer -21M13 and T in primer 241.1. Four out of five individuals are the same ones changing as in Polymorphism 6. Amino acid change occurs in edited all sequence from Y to N. There are no enzymes to cleave near this possible polymorphism.

Polymorphism 8: Frequency of base change C to T at 1142 of entire expressed sequence is 64% C and 36% T

as shown in 4 primers. No amino acid change occurs at this possible polymorphism site. Restriction enzyme cleavage site= t/cga, possible enzymes are TagI and TthHB8I. There are no other areas where this site occurs.

Polymorphism 9: Frequency of base change C to A at 1224 of entire expressed sequence is 64% C and 36% A as shown in 4 primers. The same individuals have the change from the expressed sequence as in Polymorphism 8. An amino acid change occurs in edited and edited all sequences from L to M. Restriction enzyme cleavage site=gcagc, possible enzymes are BbI and Bst71I. Other areas that this enzyme cuts are 258 and 368.

Polymorphism 10: Frequency of base change G to A at 1262 of entire expressed sequence is 57% G and 43% A. Six out of seven exhibit A. No amino acid change occurs at this possible polymorphism site. Restriction enzyme cleavage site= cctc, possible enzyme is MnlI. The enzyme cleaves the complementary DNA but not the hUCP2.GenBank sequence at this site. Other areas that this enzyme cuts are 97, 231, 278, 357, 522, 536, 594, 609, 624, 639, 695, 796, 824, 891.

*Amplification of human UCP2 exons containing polymorphisms.*

The following primers are designed to amplify hUCP2 exons 4, 6, 7 and 8 from genomic DNA. Common amino acid variants are present in exons 4, 6, and 8: A55V is in exon 4, N190S is in exon 6, and L294M is in exon 8. N190S is expected to alter a PvuII site. It is expected that Ser 190 will cut and Asn190 will not cut with PvuII. Primers for exon 7 have also been designed.

Results:

Exon 4: Both primer pairs 1 and 2 produce single bands of the expected size.

Exon 6: Primer pair 2 (3219 - 3399) produces a single band of the expected size. Primer pair 1 (3147- 3416) did not work.

Exon 7: Primer pair 1 (4316-4594) produces a single band of the expected size. Primer pair 2 (4284 - 4598) did not work.

Exon 8: Primer pairs 1 and 2 both produce multiple bands, including ones of the expected size.

PCR conditions: Clontech KlenTaq plus polymerase is used. PCR conditions were 94°C 1 min, 94°C 30 sec, 66°C 30 sec, 68°C 30 sec, repeat steps 2-4 29-times, 4°C indefinite. Reaction volume was 25 µl in an MJ research PTC-200.

=====

amplifies bp 1794-2155 of hUCP2g (Meudon)

hUCP2g.e4f1

Forward Oligo: 5'> TGTCTACTCT GTTCCCTCCC C <3'

>> Length: 21

>> Melting Temperature: 60.5

>> Max AT Run Length: 2

>> GC Clamp Strength: 57

>> % GC Content: 57

>> Hairpin Stem: 1

>> Primer Dimer: 4

>> No Palindromes.

hUCP2g.e4r1

Reverse Oligo: 5'> GGCCTACACC CTTGCTCC <3'

>> Length: 18

>> Melting Temperature: 60.6

>> Max AT Run Length: 2

>> GC Clamp Strength: 45

>> % GC Content: 67

>> Hairpin Stem: 2

>> Primer Dimer: 6

>> Longest palindrome has 4 bases (bases 1 to 4).

>> Product Size: 362

>> Annealing Temperature: 60.7 (Try T = 55.5)

=====

amplifies bp 1858-2281 of hUCP2g Meudon

hUCP2g.e4f2

Forward Oligo: 5'> TGA CTGGAGG TGGGAAGG <3'

>> Length: 18

>> Melting Temperature: 59.6

>> Max AT Run Length: 2

>> GC Clamp Strength: 47

>> % GC Content: 61

>> Hairpin Stem: 1

>> Primer Dimer: 4

>> No Palindromes.

hUCP2g.e4r2

Reverse Oligo: 5'> TATGTGGAGG ACCAGGGC <3'

>> Length: 18

>> Melting Temperature: 59.4

>> Max AT Run Length: 3

>> GC Clamp Strength: 60

SUBSTITUTE SHEET (RULE 26)



```
>> % GC Content: 61
>> Hairpin Stem: 2
>> Primer Dimer: 6
>> No Palindromes.
5 >> Product Size: 424
>> Annealing Temperature: 59.9 (Try T = 54.4)
=====
amplifies bp 3147-3416 of hUCP2g, Meudon
hUCP2g.e6f1
10 Forward Oligo: 5'> CAAGACCATT GCCCGAAG <3'
>> Length: 18
>> Melting Temperature: 60.2
>> Max AT Run Length: 3
>> GC Clamp Strength: 52
15 >> % GC Content: 56
>> Hairpin Stem: 2
>> Primer Dimer: 6
>> No Palindromes.
hUCP2g.e6r1
20 Reverse Oligo: 5'> ATGGGGGAAG GGTGAGAC <3'
>> Length: 18
>> Melting Temperature: 59.7
>> Max AT Run Length: 2
25 >> GC Clamp Strength: 14
>> % GC Content: 61
>> Hairpin Stem: 2
>> Primer Dimer: 4
>> No Palindromes.
30 >> Product Size: 269
>> Annealing Temperature: 59.2 (Try T = 54.7)
=====
amplifies bp 3219-3399 of hUCP2g, Meudon
hUCP2g.e6f2
35 Forward Oligo: 5'> TTCCTCCTCC CCGATACTC <3'
>> Length: 19
>> Melting Temperature: 59.0
>> Max AT Run Length: 3
>> GC Clamp Strength: 8
>> % GC Content: 58
>> Hairpin Stem: 2
>> Primer Dimer: 4
```

```

>> No Palindromes.
hUCP2g.e6r2
Reverse Oligo:  5'>   CCCAGCACCG TCTACCTC   <3'
>> Length:      18
5  >> Melting Temperature:    59.2
>> Max AT Run Length:    2
>> GC Clamp Strength:    21
>> % GC Content:    67
10 >> Hairpin Stem:    2
>> Primer Dimer:    4
>> No Palindromes.
>> Product Size:    180
>> Annealing Temperature:    58.1   (Try T =  54.0)
=====
15 amplifies bp 4840-5338 of hUCP2g, Meudon
hUCP2g.e8f1
Forward Oligo:  5'>   AAGTGGGAGG TGGAGGTG   <3'
>> Length:      18
>> Melting Temperature:    58.4
20 >> Max AT Run Length:    2
>> GC Clamp Strength:    29
>> % GC Content:    61
>> Hairpin Stem:    0
>> Primer Dimer:    4
25 >> No Palindromes.
hUCP2g.e8r1
Reverse Oligo:  5'>   AGCTACAAGA GAGGAGGAGA CG   <3'
>> Length:      22
>> Melting Temperature:    59.3
30 >> Max AT Run Length:    2
>> GC Clamp Strength:    30
>> % GC Content:    55
>> Hairpin Stem:    1
>> Primer Dimer:    4
35 >> Longest palindrome has 4 bases (bases 1 to 4).
>> Product Size:    520
>> Annealing Temperature:    59.5   (Try T =  53.4)
=====
amplifies bp 4831-5270 of hUCP2g, Meudon
hUCP2g.e8f2
Forward Oligo:  5'>   CAGTGAGGGA AGTGGGAGG   <3'
>> Length:      19

```

SUBSTITUTE SHEET (RULE 26)

Reverse Oligo: 5' > GGGGCAGGAC GAAGATTC <3'

```
>> Length:      18
>> Melting Temperature:      60.6
>> Max AT Run Length:      3
>> GC Clamp Strength:      19
>> % GC Content:      61
>> Hairpin Stem:      2
>> Primer Dimer:      6
>> No Palindromes.
```

```
>> Product Size:      457
>> Annealing Temperature:      60.1      (Try T = 55.6)
```

=====

amplifies 4316-4594 of hUCP2g, Meudon  
hUCP2g.e7f1

=====

Forward Oligo: 5' > CTTGCCTGCT CCTCCTTG <3'

```
>> Length:      18
>> Melting Temperature:      59.7
>> Max AT Run Length:      2
>> GC Clamp Strength:      35
>> % GC Content:      61
>> Hairpin Stem:      1
>> Primer Dimer:      6
>> No Palindromes.
```

Reverse Oligo: 5'> GGTGGTTCTC TCCCACCC <3'

```
>> Length:      18
>> Melting Temperature:      60.3
>> Max AT Run Length:      2'
>> GC Clamp Strength:      57
>> % GC Content:      67
>> Hairpin Stem:      3
>> Primer Dimer:      6
>> No Palindromes.
>> Product Size:      278
```

50

```

>> Annealing Temperature:    60.4    (Try T =  54.7)
=====
amplifies 4284-4598 of hUCP2g, Meudon
hUCP2g.e7f2
5 Forward Oligo:  5'>    CCTTCTGGAA TGATGGGTG    <3'
>> Length:      19
>> Melting Temperature:      58.9
>> Max AT Run Length:        3
>> GC Clamp Strength:        45
10 >> % GC Content:           53
>> Hairpin Stem:             2
>> Primer Dimer:             6
>> No Palindromes.
hUCP2g.e7r2
15 Reverse Oligo:  5'>    GAACTGGGTG GGGAGGAC    <3'
>> Length:      18
>> Melting Temperature:      60.3
>> Max AT Run Length:        2
>> GC Clamp Strength:        26
20 >> % GC Content:           67
>> Hairpin Stem:             2
>> Primer Dimer:             6
>> No Palindromes.
>> Product Size:           265
25 >> Annealing Temperature:    60.3    (Try T =  53.9)
=====

```

EXAMPLE X

## Immune Cell Regulation of UCP2

30 (A) A tumor mast cell line, the Rat Basophilic Leukemia cell (RBL), is a model system for studies of immune system signal transduction. In the RBL cell, binding of an antigen to the IgE receptor on the cell surface results in the activation of phospholipase

35 C-gamma (PLCγ). This event leads to the activation of protein kinase C (PKC) and an increase in intracellular calcium which are required for

SUBSTITUTE SHEET (RULE 26)

degranulation and gene expression in RBL cells. Several drugs can mimic the intracellular events and degranulation occurring in this cell type in response to antigen activation of the IgE receptor. The inophore Ionomycin causes the release of calcium from intracellular calcium stores and the phorbol ester PMA is able to directly activate PKC. Both of these drugs are required to achieve maximal activation of RBL cells. In order to determine the expression level of UCP2 in RBL cells, these cells were treated with Ionomycin (5mM), PMA (5mM) or both drugs. A Northern blot of the RNA from these cells revealed a ~4 fold induction of UCP2 expression in treated cells compared to untreated control cells. Addition of Ionomycin, PMA or both drugs simultaneously resulted in the same ~4 fold increase in UCP2 expression. Since PMA and Ionomycin treatment do not cause an additive or synergistic expression of UCP2, maximal activation of RBL cells may not be required for UCP2 gene expression.

(B) Bacterial lipopolysaccharide (LPS) regulates UCP2 mRNA in macrophages. Bacterial LPS is known to induce many of the features of infection, including fever. Thioglycolate elicited mouse macrophage (peritoneal exudate cells, PEC) have been studied for regulation of UCP2 mRNA by LPS. It has been shown that PEC express UCP2 and that LPS decreases expression by several fold over a 24 hour time course.

A study has also be made of UCP2 expression in cultured mouse macrophage RAW cells. These cells express UCP2. Addition of 100 ng/ml LPS in 5% calf serum decreases UCP2 mRNA levels by several fold after 24 hours. Lower doses of LPS are less effective, while shorter times produce smaller decreases. LPS seems to be more effective at decreasing UCP2 in cells maintained in 0.5% serum than in 5% serum. The LPS mediated decrease of UCP2 is not affected by indomethacin or by PGE2. All fo the macrophage northernns are normalized for actin expression and have been repeated at least once. These data show that LPS, which is known to influence immune function and to promote fever, can also influence immune function and to promote fever, can also influence UCP2 mRNA levels. The results are consistent with a role of UCP2 in immune function.

#### EXAMPLE XI

##### UCP2 Promoter Activity

A 246 bp putative promoter region for mouse UCP2 was cloned upstream of the chloramphenicol acetyl transferase gene in either the "sense" (also called "+" or "forward" orientation) or the "antisense" (or "-" or "reverse" orientation). When transfected into differentiated mouse HIB-1B adipocytes CAT activity (conversion of substrate into product) is observed for

the "sense" ("+") orientation, but not for the  
"antisense" ("-") orientation. (See Fig. 11).

### EXAMPLE XII

#### The Linkage Study

##### Experimental Details:

Sample. A total of 640 individuals (299 males and 341 females) from 155 pedigrees were available for the present study. These were randomly ascertained families of French descent living in the Québec city area and recruited to participate in the Québec Family Study, a population-based study of the genetics of physiological fitness and body composition. The age of individuals in the sample ranges from 18 to 94 years.

Phenotypes. BMI was obtained from height and weight measurements (BMI=weight in kg/height in m). FM, fat-free mass and %FAT were determined from body density measurements obtained by underwater weighing using the conversion factor of Siri (Adv. Biol. Med. Phys. 4:239 (1976)). RMR was determined by indirect calorimetry measurements using an open-circuit indirect calorimeter with the ventilated hood technique (Derioz et al, J. Clin. Invest. 93:838 (1994)). Measurements were taken in the morning after an overnight fast, while subjects sat quietly in a semireclined position

for the 30 minute measurement period. The last 10 minutes were kept for calculation of the RMR. The  $VO_2$  and  $VCO_2$  data were converted into energy as recommended by Weir (J. Physiol. (Lond) 109:1 (1949)). The phenotypes were adjusted by sex, for age and age by regression procedures and RMR was further adjusted for FM and FFM. The residuals from the regressions were used for linkage analysis.

*DNA typing.* Genomic DNA was prepared from permanent lymphoblastoid cells by the proteinase K and phenol/chloroform technique. DNA was dialysed four times against TE buffer (10 mM Tris, 1 mM EDTA pH 8.0) for 6 hours at 4°C and ethanol precipitated.

Amplification (EasyCycler, Ericomp, San Diego, CA) was done in 96 wells microtiter plaques using 250 ng of genomic DNA 0.1 pmoles (D11S1321, D11S916) or 0.25 pmoles (D11S911) of the forward primer coupled to the infrared tag IRD41 (Licor) and, respectively, 0.1 or 0.4 pmoles of the reverse primer, 125  $\mu$ M dNTP's, and 0.3 U Taq polymerase (Perkin Elmer, Roche Molecular Systems, Branchburg, NJ) in PCR buffer (100 mM Tris-HCl pH 8.3, 15 mM  $MgCl_2$ , 0.5 M KCl, 0.01% gelatin) for a final volume of 10  $\mu$ l. PCR cycles consisted of 1 cycle at 93°C for 5 min., 10 cycles at 94°C for 20 sec, 57°C for 60 sec, and 24 cycles at 94°C for 20 sec., 52°C for 60 sec., except for D11S911 for which the first annealing temperature was set at 55°C, PCR



products were analyzed on automatic DNA sequencer (Li-Cor) using 18 cm glass plates. Typing was done assisted by computer (One Dscan, Scannalytics).

5        *Linkage analysis.* Relative-pair based methods were used to test for linkage between the phenotypes and the marker loci. In the presence of linkage between a marker locus and a quantitative trait (Y), relative pairs sharing a greater proportion of alleles  
10       identical by descent ( $\pi$ ) as the marker locus tend to have more similar phenotypes than pairs who share fewer alleles. Thus, under the hypothesis of linkage, a negative relationship is expected between  $\pi$  and the within pair variance. The sib-pair linkage method  
15       described by Haseman and Elston is the most widely used method to investigate linkage between a quantitative phenotype and a marker locus. This method has been extended to other types of relative pairs. Tests of linkage which combine the information  
20       from different types of relative pairs were developed and these tests have been shown to be more powerful than the Haseman and Elston method based only on sib-pairs. This relative pair linkage analysis has been implemented in the program RELPAL which considers  
25       the following five types of relative pairs: sibling, half-sibling, grandparent-grandchild, avuncular and first degree cousins. For each relative pair type, the statistic for testing linkage is obtained by

dividing the estimated regression coefficient ( $\beta$ ) by its standard error. Because the number ( $n$ ) of relative pairs could vary among the different types of relatives depending on the complexity of the pedigrees, the contribution of each type of relative pairs needs to be weighted in the overall linkage statistic which combine information from all relative pairs. The linkage test implemented in RELPAL is:

$$T = \frac{c^T \beta}{\sqrt{c^T \text{Var}(\beta) c}}$$

where  $\beta$  is a vector containing the  $\beta$ 's for each of the 5 types of relatives and  $c^T$  is a weighing vector based on  $n$  and the variance of  $\pi$  and equal to:  $(\text{Var}(\pi_s)n_s, \text{Var}(\pi_h)n_h, \text{Var}(\pi_g)n_g, \text{Var}(\pi_a)n_a, \text{Var}(\pi_c)n_c)$ , where subscripts  $s, h, g, a$  and  $c$  stand for siblings, half-siblings, grandparent-grandchild, avuncular and first degree cousins, respectively. In the Québec Family Study, since there are no half-sibs, only sibling, avuncular, grandparental and cousin pairs were used in the relative-pair linkage analysis.

#### Results:

A study was undertaken to investigate the potential linkage relationships between three microsatellite markers which encompass the UCP2 gene location on 11q13 with resting metabolic rate (RMR), body mass index (BMI), percentage body fat (%FAT) and fat mass (FM) in 640 individuals from 155 randomly

ascertained pedigrees from the Québec Family Study. Using a linkage analysis strategy based on sibling, avuncular, grandparental and cousin pairs, strong evidence of linkage was found between the marker D11S911 ( $p=0.000002$ ) and RMR with more moderate evidence for D11S916 ( $p=0.006$ ) and D11S1321 ( $p=0.02$ ). Suggestive evidence of linkage was also observed between D11S1321 and %FAT ( $p=0.04$ ) and FM ( $p=0.02$ ). It is concluded that the three markers encompassing the UCP2 locus and spanning a 5 cM region on 11q13 are linked to resting energy expenditure in adult humans.

The mouse *ucp2* gene was recently mapped to chromosome 7, closely linked to the *tubby* mutation, a mutation known to be responsible for the adult-onset obesity in this mouse model. Furthermore, the UCP2 mRNA level was found to be higher in the A/J mouse strain, which is resistant to diet-induced obesity, than in the obesity prone C5/BL/6J mouse. The evidence accumulated thus far on animal models indicates that the UCP2 gene plays a role in the development of obesity because of its role in energy metabolism. The human UCP2 gene has been mapped to chromosome 11q13 at a location distinct from *tubby* (11p15.1) but in the same chromosomal location as the Bardet-Biedl Syndrome locus (Online Mendelian Inheritance in Man, OMTM, Center for Medical Genetics, Johns Hopkins Univ., (Baltimore, MD) and National Center for Biotechnology Information, National Library

of Medicine (Bethesda, MD) (1986), WWW URL:  
<http://www3.ncbi.nlm.nih.gov/omim/>), a Mendelian  
Syndrome exhibiting obesity as one of its clinical  
features. UCP2 is also in the proximity (about 15 cM)  
of a locus (11q21-q22) recently uncovered through a  
genome-wide search and found to be linked to percent  
body fat in Pima Indians. To test the hypothesis that  
markers around the UCP2 gene may exhibit a linkage  
relationship with metabolic rate and body fat  
phenotypes, three markers (D11S916, D11S1321 and  
D11S911) were typed on 640 individuals from 155  
pedigrees from the Québec Family Study. Linkage  
studies were undertaken with RMR, BMI, %FAT and FM  
using four types of relatives. RMR was adjusted for  
the effects of age, sex, FM and fat-free mass, whereas  
BMI, %FAT and FM were adjusted only for age and sex  
effects.

Table 3 presents the linkage results with the  
number of relative pairs available in each case.  
Strong evidence of linkage was observed between the  
D11S911 and RMR while more moderate evidence for  
linkage was found for the other two markers. As there  
were few pairs of first degree cousins, the same  
analysis was repeated after removing these pairs. In  
the latter case, the p-value for linkage between  
D11S911 and RMR (n=212 pairs) was reduced to p=0.001,  
while the one between D11S916 and RMR (n=279 pairs)  
went from p=0.006 to p=0.02. Evidence of linkage

between D11S1321 and %FAT ( $p=0.04$ ) and FM ( $p=0.02$ ) was also found. No linkage was observed between the three markers and fat-free mass in this population.

Overall, these results indicate that the UCP2 gene which is encoded within the 5 cM span covered by these markers, plays a role in determining resting energy expenditure in humans.

Table 3. Relative-pair linkage analysis of resting metabolic rate and body fat variables with markers encompassing the UCP2 gene in the Québec Family Study

MARKER	HZ	cM	RMR	BMI	%FAT	FM
D11S916	0.72	78	0.006 (301)	0.27 (415)	0.50 (304)	0.11 (304)
D11S1321	0.64	79	0.02 (380)	0.36 (537)	0.04 (383)	0.02 (383)
D11S911	0.85	83	0.000002 (240)	0.38 (324)	0.23 (243)	0.26 (243)

Based on four different types of relative pairs: siblings (165-275 pairs); avuncular (47-134 pairs); grandparental (45-94 pairs); first degree cousins (22-34 pairs). The entries are the p-values with total number of relative pairs given in parentheses. HZ: heterozygosity; cM; distance in centimorgan.

\* \* \*

All documents cited herein are incorporated in their entirety by references.

What is claimed is:

1. A method of treating a disease or disorder associated with diminished UCP2 expression or activity comprising administering to an individual in need of such treatment an amount of an agent that enhances UCP2 expression or activity sufficient to effect said treatment.
2. The method according to claim 1 wherein said disease or disorder is obesity, diabetes, syndrome X, or hypothermia.
3. The method according to claim 1 wherein said disease or disorder is hyperinsulinemia or glucose intolerance.
4. The method according to claim 1 wherein said agent is an expression construct comprising a UCP2 encoding sequence or portion thereof that encodes functional UCP2.
5. A method of treating a disease or disorder associated with elevated UCP2 expression or activity comprising administering to an individual in need of such treatment an amount of an agent that inhibits UCP2 expression or activity sufficient to effect said treatment.

6. The method according to claim 5 wherein said disease or disorder is wasting, anorexia, inflammation, cachexia, fever, or hyperthermia.
7. The method according to claim 5 wherein said agent is an oligonucleotide that specifically hybridizes to UCP2 mRNA or an antibody, or binding fragment thereof, that is specific for UCP2.
8. A method of screening a compound for its ability to modulate UCP2 expression or activity comprising contacting a cell that expresses UCP2 with said compound under conditions such that modulation of UCP2 expression or activity can occur and measuring UCP2 expression or activity in the presence and absence of said compound and determining whether modulation is effected.
9. Isolated mammalian UCP2, or fragment thereof of at least 5 amino acids.
10. The UCP2 according to claim 9 wherein said UCP2 is human UCP2.
11. The UCP2 according to claim 10 wherein said UCP2 has the sequence shown in Figure 1A.



12. An isolated nucleic acid sequence encoding mammalian UCP2, or portion thereof at least 15 nucleotides in length.

13. The nucleic acid according to claim 12 wherein said nucleic acid encodes human UCP2.

14. The nucleic acid, according to claim 13 wherein said human UCP2 has the sequence shown in Figure 1A.

15. A recombinant molecule comprising a vector and the nucleic acid according to claim 12.

16. A host cell comprising the recombinant molecule according to claim 15.

17. A method of making UCP2 comprising incubating the host cell according to claim 16 under conditions such that said nucleic acid is expressed and said UCP2 is thereby produced.

18. An antibody specific for UCP2 or antigen binding fragment thereof.

19. An antisense construct comprising a vector and a nucleic acid operably linked to a promoter, wherein said nucleic acid is oriented with respect to the

promoter such that the transcript of the nucleic acid selectively hybridizes to UCP2 mRNA.

20. A host cell comprising the construct according to claim 19.

21. The cell according to claim 20 wherein said cell is a mammalian cell.

Fig. 1A.

DEFINITION hUCP2, 309 amino acids.  
ORIGIN

```

1  MVGFKATDVP PTATVKFLGA GTAACIADLI TFPLDTAKVR LQIQGESQGP
51  VRATASAQYR GVMGTILTMV RTEGPRSLYN GLVAGLQRM SFASVRIGLY
101 DSVKQFYFKG SEHASIGSRL LAGSTTGALA VAVAQPTDVV KVRFQAQARA
151 GGGRRYQSTV NAYKTIAREE GFRGLWKGS PNVARNAIVN CAELVTYDLI
201 KDALLKANLM TDDLPCHFIS AFGAGFCTTV IASPVDVVKT RYMNSALGOY
251 SSAGHCALTM LQKEGPRAFY KGFMPSFLRL GSWNVVMFVT YEQLKRALMA
301 ACTSREAPF

```


Fig 1B.

DEFINITION hUCP2 coding sequence, 930 bases including the stop  
codon.  
ORIGIN

```

1  ATGGTTGGGT TCAAGGCCAC AGATGTGCCC CTTACTGCCA CTGTGAAGTT
51  TCTTGGGGCT GGCACAGCTG CCTGCATCGC AGATCTCATC ACCTTTCCTC
101 TGGATACTGC TAAAGTCCGG TTACAGATCC AAGGAGAAAG TCAGGGGCCA
151 GTGCGCGCTA CAGCCAGCGC CCAGTACCGC GGTGTGATGG GCACCATCTT
201 GACCATGGTG CGTACTGAGG GCCCCCGAAG CCTCTACAAT GGGCTGGTTG
251 CCGGCCTGCA GCGCCAAATG AGCTTTGCCCT CTGTCCGCAT CGGCCGTGAT
301 GATTCTGTCA AACAGTTCTA CACCAAGGGC TCTGAGCATG CCAGCATTTG
351 GAGCCGCCCTC CTAGCAGGCA GCACCACAGG TGCCCTGGCT GTGGCTGTGG
401 CCCAGCCCCAC GGATGTGGTA AAGTCCGAT TCCAAGCTCA GGCCCGGGCT
451 GGAGGTGGTC GGAGATACCA AAGCACCGTC AATGCCCTACA AGACCATTGC
501 CCGAGAGGAA GGGTTCCGGG GCCTCTGGAA AGGACCTCTT CCAATGTTG
551 CTCGTAATGC CATTGTCAAC TGTGCTGAGC TGGTGACCTA TGACCTCATC
601 AAGGATGCCC TCCTGAAAGC CAACCTCATG ACAGATGACC TCCCTTGCCA
651 CTTCATTTCT GCCTTTGGGG CAGGCTTCTG CACCACTGTC ATCGCCTCCC
701 CTGTAGACGT GGTCAAGACG AGATACATGA ACTCTGCCCT GGGCCAGTAC
751 AGTAGCGGTG GCCACTGTGC CTTTACCATG TCCAGAAAG AGGGCCCCCG
801 AGCCTTCTAC AAAGGTTTCA TGCCCTCCTT TCTCCGCTTG GGTTCCTGGA
851 ACGTGGTGAT GTTCGTCACC TATGAGCAGC TGAACGAGC CCTCATGGCT
901 GCCTGCACTT CCGAGAGGC TCCCTTCTGA

```




sequence data not yet available

UCP2 exons (1-6)

Fig. 1C.

restriction sites: fragment size indicated (Kb) when





**Fig. 1D. (continued 1)**

.....

pSUB23t7

.....

GGATCCCTAAGGAGAAGAGCCTGGAGGAGATGAGAAGTGGGGAGGAAGGCCTGCTGCGCA  
GATCTGGTGTACAGATCCAGGCGGTGCAGATGAGCAGATGGGACCAAGTGGATGGTGATT  
TAGAGGGTGGAATCCACAGGGCTTAGTGACTGATGGCAGGTGGTTGTGGGAGGGGAGGCC  
AAGTCTAGGCCGGGGCAGGCAGGGCTCACTCTGATCTCCTCCCCGCTGCTCCTGCACTTT  
GCCCCTCACCTTGCCAAGAAGGATNGGGTTCACAAAAAAAGTTGTCCTTCNCCCCTGC

.....

pSUB5t7

.....

GGATCCCCCTAAGTTGGAGGAAAGGAAACTGGCCCAAACAAAAAGGAGAGCAGTTTTCTC  
TGCATCACATGGTAGGCCAGGAGGAGTCTAATGCCCCAGAGTTTACTCTCAGCCCCCAA  
ATCACCTAGCTAAATGTTACCTTATCTAAGAAGTCCTTAGGTTTTTGGGGTTTGT  
TTTTTTTTGAGACAAGGTCTCACTCTCTCACCAGACTGGAGCACAGTGGCACAATCAC  
AGCTCACTGCAGCCTCAACCTCCTGGGCTCAAGCAATCGTCCCAAGTAGCTGGGACTATA  
GGCCTGCACCACCATGTTCACTAATTTATTTTATTTATTTTTAAAAGGTCTCATTAT  
TTGCCTGCTGGTCTGAATCCTGGTTCAACATCTCCACC

.....

ex12

.....

GAGCTCACATCGGTTTTCCCTCATGAGGCCACTTGGAGTCTTGCTGAGGGACTTGGTTCT  
ATTAGGGAAGGTGAGTTTGGGGATGGTGAGCAGGGAGGGCCTGGGGACATTGTGGCTAAT  
GGGGCTTTTCTCCTCTTGGCTTAGATTCCGGCAGAGTTCCTCTATCTCGTCTTGTTGCTG  
ATTAAAGGTGCCCTGTCTCCAGTTTTTCTCCATCTCCTGGGACGTAGCAGGAAATCAGC  
ATCATGGTTGGGTTCAAGGCCACAGATGTGCCCCCTACTGCCACTGTGAAGTTTCTTGGG  
GCTGGCACAGCTGCCTGCATCGCAGATCTCATCACCTTTCCTCTGGATACTGCTAAAGTC  
CGGTTACAGGTGAGGGGATGAAGCCTGGGARTCTTGATGGTGTCTACTCTGTTCCCTCCC  
CAAAGACACAGACCCCTCAAGGGCCAGTGTGGAGCATCGAGATGACTGGAGGTGGGAA  
GGGCAACATGCTTATCCCTGTAGCTACCCTGCTTGGCCTTGACAGATCCAAGGAGAAAGT  
CAGGGGCCAGTGCGCGCTACAGCCAGCGCCAGTACCGCGGTGTGATGGGCACCATTCTG  
ACCATGGTGCGTACTGAGGGCCCCCGAAGCCTCTACAATGGGCTGGTTGCCGGCCTGCAG  
CGCCAAATGAGCTTTGCCTCTGTCCGCATCGGCCTGTATGATTCTGTCAAACAGTTCTAC  
ACCAAGGGCTCTGAGCGTGAGTATGGAGCAAGGGGTGTAGGCCCTTGGCCCTTTTTTCTC  
AGTGATGATTGATCTTAGTTCATTACGCCATATAGTTTTTTAGGCCCCACGATCCCTAGG  
AAGATCAGGGGGAACAGAGAACTGGAAGGGGCCCTGGTCCTCCACATAGTTCCTAAGCAC  
CTGGGCTATACCAGGCTCTGAGCAGGGCGTCATCCCATCACAGTCTTCAACACCACCTTG  
GGAGTAGGTAGTATCATCCAGTGTTATAGAAGAAGAGACTGAGGTGGGAAGGCAGTGGG  
TAGAGTGGGGACTTGGCCAGGGGCACACAGTAGAGAGCCAGAAAACACACAGTAGAGAGC  
CAGGACACTCGTCTCTAAGGCCAGCGTCTTCCCTTTACCTCCTTAGTATGCCATGCCA  
ACCCTCCATTTTACACATGACGAAACAGAGCCCCAAACAAAAGTTGTCTTTCCAGATC  
ACATGGCAGGAAGAAGTAAAGCTGACCTGAGATCCCAAGTCTTAGGAATCCAGTCCTCA  
GAAAGCCACTTCTCTCTGAGCCTTGGTTTTACATTTGTCAGATGGAAATGATTGTGATT  
TCTCAGGGCTGTTGAGCAGGTAAATGAAAATGTTTTATGAAAGAAAGCACCAAGTTTCAT  
TTTGGTCTTAGCCCTTGCTATGTCCCTAGCAAGAAGTAGATATTCATAGGGATATTTTGT  
TTGATGTGAGGAGTCTTACAGCAAGAGCTTGTAGAAGGCCAAAAGCTTCTGGATTCTAA  
TCCAAAAGCAGGAGATGACAGTGACAGGGTGGTTTTGGTGAGGAGAGATGAGGTAGAAA  
ATGAGTGCAAGCCCGCTGGCCACTGACCCCATGGCTCGCCACAGATGCCAGCATTGGGG  
CNTNCCNTTGG

**Fig. 1D. (continued 2)**

.....

ex34

.....

CCCGNTGNGCTGTGGCTGTGGCCCAGCCCACGGATGTGGTAAAGGTCCGATTCCAAGCTC  
AGGCCCCGGGCTGGAGGTGGTCGGAGATACCAAAGCACCGTCAATGCCTACAAGACCATTG  
CCCCGAGGGAAGGGTTCCGGGGCCCTCTGGAAAGGTGTGTACCAGTTGTTTTCCCTTCCCC  
TTTTCTCCTCCCCGATACTCTGGTCTCACCCAGGATCTTCTCCTCCTACAGGGACCTC  
TCCCAATGTTGCTCGTAATGCCATTGTCAACTGTGCTGAGCTGGTGACCTATGACCTCAT  
CAAGGATGCCCTCCTGAAAGCCAACCTCATGACAGGTGAGTCATGAGGTAGACGGTGCTG  
GGTCTCACCTTCCCCCATGCCAGGARCAAGGTGCGGGGGTCTAGCTGACACCAGAAGACC  
ACATCTTTTCATCCTATTTGCCCTTTGCAGGGAGAGTAAGATATCTCTTACTTGCCATAT  
TGAAGCCAATTGGGATGAAGCTCCCACTTTGCACATTGAGGAACTGAGGCTAGATTGGCA  
AAATGACTCTTTCAGGTCCTCAGAAGATGTCTCAGCTGGAGTCCCTGTCTGTTTTGTTT  
TTTTGTTGTTTGTGTTTTGTTTTTGTGTTTTTGTGAGATAGANTCTCACTCTGTTACCCGTGTAAT  
CTCAGCTCACTGCAACCTTCTCCTCCTGGGTTCAAGCAATCCTGTGCCTCACCTCCC

.....

ex56

.....

GTCATGGAGCTTGACTTCCGGATTAAGCATCTTCCACTTCAGCCTCTNGAGTAGCTGGGC  
TGACTACAGGCATGCACACTGTGCCTGGCTAATTTTNGTATTTTGTAGAGACAGGGTTTT  
TGCCATGTTACCCAGTCTGGTCTTGAACCTCCTGGGCTCAAGTGATCCACCCACNTCGGCC  
TCCAAAAGAAGTCCTGGATTACAGGCATGAGACATTGTGCCCAGCCTCTCTGTCTCTTTA  
AAATCATGAAAACCTCGTAGCTACTTAAGTAATTCTCCTGCCTTCTGGAATGATGGGTGAA  
GATCTTGACTGCCTTGCTCCTCCTTGGCAGATGACCTCCCTTGCCACTTCACTTCT  
GCCTTTGGGGCAGGCTTCTGCACCACTGTATCGCCTCCCTGTAGACGTGGTCAAGACG  
AGATACATGAACTCTGCCCTGGGCCAGTACAGTAGCGCTGGCCACTGTGCCCTTACCATG  
CTCCAGAAGGAGGGCCCCGAGCCTTCTACAAAGGGTGAGCCTCTGGTCTCCCCACCCAG  
TTCAGGCNTCTTGGCTATGCATGTCTATTATGGGTGGGAGAGAACCACCTGGAAGTGAGT  
AGCAGCCAAGTGTGACTATTTCTGATCCTGGTCGTGGCATTTCACCAGCATTACCTATC  
CCCTTAATTCCTTCCCTCCCAGAATTGCTACCATCACTGTTTATTAGGTGTTAAATGGAGA  
CTCAAAGGGAATTCATGCTTATAGCCAAGCAGCTGTGAGCTCAGTTCATTGAGTCCTCCC  
AGCCTCCTTTGGGACAGAGCAACTGGGTTGGATTGAATACCAGGCCAGTGAGGGAAGTG  
GGAGGTGGAGGTGCCCCCATGACCTGTGATTTTTCTCCTCTAGGTTTCATGCCCTCCTTTC  
TCCGCTTGGGTTCTGGAACGTGGTGATGTTCTGTCACCTATGAGCAGCTGAAACGAGCCC  
TCATGGCTGCCTGCACTTCCCGAGAGGCTCCCTTCTGAGCCTCTCCTGCTGCTGACCTGA  
TCACCTCTGGCTTTGTCTCTAGCCGGGCCATGCTTTCTTTTCTTCTTCTTCTTCTTCC  
CTCCTTCCCTTCTCTCCTTCCCTCTTTCCCCACCTCTTCTTCCGCTCCTTTACCTACCA  
CCTTCCCTCTTCTACATTCTCATCTACTCATTGTCTCAGTGCTGGTGGAGTTGACATTT  
GACAGTGTGGGAGGCCTCGTACCAGCCAGGATCCCAAGCGTCCCGTCCCTTGAAAGTTC  
AGCCAGAATCTTCGTCTGCCCCGACAGCCCAGCCTAGCCCACTTGTCATCCATAAAGC  
AAGCTCAACCTTGGCGTCTCCTCCCTCTCTTGTAGCTCTTACCAGAGGTCTTGGTCCAAT  
GGCCTTTTTTGGTACCTGGTGGGAGGGAGGAACCACTGACTTTGAAATGGGTGTGAT  
CCACCTTCCACCTCCAGCATCCAATCTGAAGCCCGTGTAGGTCATCTGGTCCATTTCTCT  
CTAGATCCAGGCCCTGTTACTAACATGGGGAGTGAGGAGCCACCTGAGAGACAGCAGTG  
CCTCCCCTTCTTTGCCGGGCCACTTGAGCTC

**Fig. 1E.**

LOCUS hUCP2.genebank 1612 bp  
 DEFINITION hUCP2.genebank, 1612 bases.

```

1  TGA CTGTCCA CGCTCGCCCCG GCTCGTCCGA CGCGCCCTCC GCCAGCCGAC
51 AACACAGCCG CACGCACTGC CGTGTTCCTCC CTGCGGCTCG GACACATAGT
101 ATGACCATTA GGTGTTTCGTC TCCCACCCAT TTTCTATGGA AAACCAAGGG
151 GATCGGGCCA TGATAACCAC TGGCAGCTTT GAAGAACGGG ACACCTTTAG
201 AGAAGCTTGA TCTTGGAGGC CTCACCGTGA GACCTTACAA AGCCGGATTC
251 CGGCGAGTTC CTCTATCTCG TCTTGTGCT GATTAAAGGT GCCCCTGTCT
301 CCAGTTTTC TCCATCTCCT GGGACGTAGC AGGAAATCAG CATCATGGTT
351 GGGTTCAAGG CCACAGATGT GCCCCCTACT GCCACTGTGA AGTTTCTTGG
401 GGCTGGCACA GCTGCCTGCA TCGCAGATCT CATCACCTTT CCTCTGGATA
451 CTGCTAAAGT CCGGTTACAG ATCCAAGGAG AAAGTCAGGG GCCAGTGCGC
501 GCTACAGCCA GCGCCCAGTA CCGCGGTGTG ATGGGCACCA TTCTGACCAT
551 GGTGCGTACT GAGGGCCCCC GAAGCCTCTA CAATGGGCTG GTTGCCGGCC
601 TGCAGCGCCA AATGAGCTTT GCCTCTGTCC GCATCGGCCT GTATGATTCT
651 GTCAAACAGT TCTACACCAA GGGCTCTGAG CATGCCAGCA TTGGGAGCCG
701 CCTCCTAGCA GGCAGCACCA CAGGTGCCCT GGCTGTGGCT GTGGCCCAGC
751 CCACGGATGT GGTAAGGTC CGATTCCAAG CTCAGGCCCCG GGCTGGAGGT
801 GGTCGGAGAT ACCAAAGCAC CGTCAATGCC TACAAGACCA TTGCCCGAGA
851 GGAAGGGTTC CGGGGCCTCT GGAAAGGGAC CTCTCCCAAT GTTGCTCGTA
901 ATGCCATTGT CAACTGTGCT GAGCTGGTGA CCTATGACCT CATCAAGGAT
951 GCCCTCCTGA AAGCCAACCT CATGACAGAT GACCTCCCTT GCCACTTCAT
1001 TTCTGCCTTT GGGGCAGGCT TCTGCACCAC TGTCATCGCC TCCCCTGTAG
1051 ACGTGGTCAA GACGAGATAC ATGAACTCTG CCCTGGGCCA GTACAGTAGC
1101 GCTGGCCACT GTGCCCTTAC CATGCTCCAG AAGGAGGGGC CCCGAGCCTT
1151 CTACAAAGGG TTCATGCCCT CCTTTCTCCG CTTGGGTTCC TGGAACGTGG
1201 TGATGTTTCG CACCTATGAG CAGCTGAAAC GAGCCCTCAT GGCTGCCTGC
1251 ACTTCCCGAG AGGCTCCCTT CTGAGCCTCT CCTGCTGCTG ACCTGATCAC
1301 CTCTGGCTTT GTCTCTAGCC GGGCCATGCT TTCCTTTTCT TCCTTCTTTC
1351 TCTTCCCTCC TTCCCTTCTC TCCTTCCCTC TTTCCCCACT TCTTCTTCC
1401 GGGCTCCTTA ACCTACCAAC CTTCCCTCTT TCAACATTCT CATCTACTCA
1451 TTGTCTCAGT GCTGGTGGAG TTGACATTTG ACAGTGTGGG AGGCCTTGTA
1501 CCAGCCAGGA TCCCAAGCGT CCCGTCCCTT GGAAAGTTCA GCCAGAATTT
1551 TTGTCCTGCC CCCGACAGCC CAGCCTAGCC CATTGTCATC CATAAAGCAA
1601 GCTCAACCTT GG

```



## hUCP2.CDSF1

Forward Oligo: 5'&gt; CATCTCCTGG GACGTAGC &lt;3'

>> Length: 18  
>> Melting Temperature: 56.6  
>> Max AT Run Length: 2  
>> GC Clamp Strength: 42  
>> % GC Content: 61  
>> Hairpin Stem: 2  
>> Primer Dimer: 6  
>> Longest palindrome has 4 bases (bases 12 to 15).

## hUCP2.CDSR1

Reverse Oligo: 5'&gt; GGAGAGAAGG GAAGGAGGG &lt;3'

>> Length: 19  
>> Melting Temperature: 60.1  
>> Max AT Run Length: 2  
>> GC Clamp Strength: 57  
>> % GC Content: 63  
>> Hairpin Stem: 0  
>> Primer Dimer: 0  
>> No Palindromes.

&gt;&gt; Product Size: 1098

&gt;&gt; Annealing Temperature: 59.4

Try T = 51.6

## Pair 2

## hUCP2.CDSF2

Forward Oligo: 5'&gt; AGGTGCCCCT GTCTCCAG &lt;3'

>> Length: 18  
>> Melting Temperature: 61.3  
>> Max AT Run Length: 1  
>> GC Clamp Strength: 32  
>> % GC Content: 67  
>> Hairpin Stem: 2  
>> Primer Dimer: 6  
>> No Palindromes.

**Fig. 1F.**

## hUCP2.CDSR2

**Fig. 1F. (continued 1)**

Reverse Oligo: 5'> TCAGGTCAGC AGCAGGAG <3'

>> Length: 18  
>> Melting Temperature: 58.7  
>> Max AT Run Length: 1  
>> GC Clamp Strength: 32  
>> % GC Content: 61  
>> Hairpin Stem: 2  
>> Primer Dimer: 6  
>> No Palindromes.

>> Product Size: 1043  
>> Annealing Temperature: 60.1                      Try T = 53.7

---

Pair 3  
hUCP2.CDSF3

Forward Oligo: 5'> CATCTCCTGG GACGTAGC <3'

>> Length: 18  
>> Melting Temperature: 56.6  
>> Max AT Run Length: 2  
>> GC Clamp Strength: 42  
>> % GC Content: 61  
>> Hairpin Stem: 2  
>> Primer Dimer: 6  
>> Longest palindrome has 4 bases (bases 12 to 15).

hUCP2.CDSR3

Reverse Oligo: 5'> AGAGAAGGGA AGGAGGGAAG <3'

>> Length: 20  
>> Melting Temperature: 58.9  
>> Max AT Run Length: 2  
>> GC Clamp Strength: 47  
>> % GC Content: 55  
>> Hairpin Stem: 0  
>> Primer Dimer: 0  
>> No Palindromes.

**Fig. 1F. (continued 2)**

>> Product Size: 1096  
>> Annealing Temperature: 59.4 Try T = 51.6

---

Pair 4  
hUCP2.CDSF4

Forward Oligo: 5' CATCTCCTGG GACGTAGC 3'

>> Length: 18  
>> Melting Temperature: 56.6  
>> Max AT Run Length: 2  
>> GC Clamp Strength: 42  
>> % GC Content: 61  
>> Hairpin Stem: 2  
>> Primer Dimer: 6  
>> Longest palindrome has 4 bases (bases 12 to 15).

hUCP2.CDSR4

Reverse Oligo: 5' AAGGAGAGAA GGGAAGGAGG 3'

>> Length: 20  
>> Melting Temperature: 58.9  
>> Max AT Run Length: 2  
>> GC Clamp Strength: 43  
>> % GC Content: 55  
>> Hairpin Stem: 0  
>> Primer Dimer: 0  
>> No Palindromes.

>> Product Size: 1100  
>> Annealing Temperature: 59.4 Try T = 51.6

---

Pair 5  
hUCP2.CDSF5

Forward Oligo: 5' AGGTGCCCCT GTCTCCAG 3'

>> Length: 18

SUBSTITUTE SHEET (RULE 26)

**Fig. 1F. (continued 3)**

>> Melting Temperature: 61.3  
>> Max AT Run Length: 1  
>> GC Clamp Strength: 32  
>> % GC Content: 67  
>> Hairpin Stem: 2  
>> Primer Dimer: 6  
>> No Palindromes.

hUCP2.CDSR5


Reverse Oligo: 5'> AGGAGAGAAG GGAAGGAGG <3'

>> Length: 19  
>> Melting Temperature: 57.4  
>> Max AT Run Length: 2  
>> GC Clamp Strength: 43  
>> % GC Content: 58  
>> Hairpin Stem: 0  
>> Primer Dimer: 0  
>> No Palindromes.

>> Product Size: 1125  
>> Annealing Temperature: 60.8

Try T = 52.4

Fig. 2.



**Fig. 3A.****Human**


H B Pl Lu L M K P



UCP2

**Mouse**

B K L Wat Bat H M1 M2



UCP1







**Fig. 3B.****Fig. 3C.**UCP2 RNA/cyclophilin RNA  
(phosphorimager units)**Fig. 4.**

Fig. 5.




**Fig. 6.**

HIB 1B cells grown in DMEM + 10% charcoal-stripped serum (*Media*) or with the addition of the thiazolidine dione BRL 49653 and the RXR $\alpha$  ligand LGD-1069 ( *TZD/RXR*). The results are the average of 2 independent experiments of 4 samples each. \*, significantly different from *Media* samples,  $p < 0.001$ .



## Map of the mouse UCP2 gene.

Clone "MMU2-L2" (insert 13.9 Kb) deposited at Pasteur Institute  
(Registration number: I-1868, 04.16.1997)

**Fig. 7.**

SUBSTITUTE SHEET (RULE 26)

**MMU2-L2: partial promoter sequence (unmapped)**

[illegible]

**promoter and coding sequence:**

[illegible]

**SUBSTITUTE SHEET (RULE 26)**


[illegible]

TcACAAtGCCcTcCCAGGATcGcTcTcCAGGGAGAGtAGAGcTGGGgCTAAAGAAcTctGccTcTgGAcTCAGAcTcTGCTAGAGAAAAGccTcTfGcTTGgTcttgGgtTTCACATTTGATAAAATGGAaCTGACCCtGAAaTTACCAAGGCTGCGAGAGTAGCGTAAATGAACACATTTATGTAGAAAGCGCAGTCTCCATTTTGTGTtAcATcCTGGTGTAAACCTTCGAAGAGTGATCTAGAGCcTCTCTCTTCCCGTGGCATGGTGGTCTGGCAAGAGCATGATGCACTTGAGTGCAGCcTCAGCcTCGGGCCACTAACCCCGCCCCAGCCcTCTTTGCCACAGATGCAGGCATCGGAGCCGCCCTCTTGGCAGGTAGCACCAACAGGTgCCCTGGCCGTGGCTGTAGCCCCAGCCTACAGATGTGGTAAAGGTCCGCTTCAGGCTCAGGCCCGGGCTGGTGGTGGTGGAGATACACAGACACTGTGCGAAGCCCTACAAGACCAtGCACGAGAGGAAGGATCCGGGCCCTCTGGAAAGGTGTGTGCTGGGTGTGGCTTTTCTCCCCCTCTCTCTCCATGCCCCTGGCTCACACAGGTCTCTCTTTTCTACAGGGACTCTCTCCAAATGTTGCCCCGTAAATGCCATTTGTAACCTGTGCTGAGCTGGTGACCTATGACCCTCAATAAGATACTCTCTCTGAAAGCCAACTCATATGACAGGTGTGTGTTAGCGCCACTTCAAGCTGGTAGGGGGCGTctAGTTGTCTAAAGACTGCATctTGTtATCTCTGCTTGCCTCGTGGTGTTTTAGAGTTTATATTTCTTACCATTCAAATCCAACAGGGTAAACCCCACTTTGCACATAAACAGCGGAGATCAACAAaAAGACTCTGAAGTCCCCCTAAGAGAGTCTAGCTGTAGTCTGTCTTTAAACAAATAGAGAcCCCTCTGcgGCAaTGGTGGGgAtGCgGctTgGGAGgctGAbAgCcTcTCCCCCTGCTTCATGGCAGatGACcTCCctTGcAcTTCAcTTcTcTtTGCCcTtTcgGGGCGGcttctGCACCACCGhCATCGCCCTCCCCCTGTGTGTGATGTcAGAGcAGATACCTGCTGCTTGGGCCCAGTACCACAGCGCAGTCAcTGTGCCCTTACCATGCTCCGGAAGGAGGGACCCcgCGCTTCTAcAaGGGgTGAGCCcCagATcCCcTCAcAGAAcTCCAGAGAGTcAGTcAGCCAAATGTcAGACAGCCCCATAGTTGCTGTGTACcTCTGTtTtTtTAGAAGTTAAATGAGCATTTCAgAcTGGGGGTGGGATGGGTGGGTcAGTGGTGGAGGGCTTGTCTAGCATGCTTGGAGACCCtTGGGTTCTGTCTCCCAACAGCAGGAGTAgGGAGCTTGGGTACTCTGGGATTTCTTGGGCATGTGGCCATGGGTGGGTGGATGGATGAGCCcTGTGTAGAAcCAGAGTGGTGGAGGACCTCCATGACTGTGTtTGTCTcTcTcTAGGTTcATGCCtTCTCTTCTCGCTGGGATCTCTGGAAcGTAGTATTTGTCAcCTATAGAGCCTTAAGAGCCCCtAATGGCTGCCtTACCAATCTCGGGAGCCTTTCTGAGCCCTCTCCATGCTGACCTGGAGCCCTGCTTCCCAAGCCCTGCTcTtTtTtTCTCATCTCTGTCCAGTCCCATTTCTTCCCATTTCTCCATGCTGACCCGATTTACTTTCCCAcCTcCTGCTGCCCTCTGTACCTCTGTACTGATGACTCAcAGTGAGGAGGCCtGACACcAGACCCTGAGCCCTCAGCCCTTTCACAGCTACAGCTAAGCCCAcATCTTTcATCTTCAATCCCCAGCCcAGCCcDAGCCcAGCTcAGCCAGCCTTcACCCATAAAGCAAGCTCAATGTGTGGTGTCTTCTTCTCATATGTTTACAGAGGTCTTGTGTCCAAAGGAcTTTtTTCAGGACcTTTGGCAGGCAGGTGAGGACCCcAgTGTCTTTTAAAAAGGgAtGTGACCCcACA TTcTGCcTGTAAcATcTgAGAGg99CTGGAnGGTcTtCCAGTCTGTTCtTcAgACCCAgTGTtGGcTCCAgAgATCAgTGGCbaCAGCCAgGGGAcACTGCAgCCtTTGTccCTCCATCCcTtTcCTGAAcACCATGTAAAGCTGTACCTCAGTACCTCGTACTCCACACACTGGCCAGGGCTGTCTATCTGcAGTGGAAgTATCCCTnCcCTCCCAcATCAACTTGTtTAGGAGA-TGTAGGGAAAcTCCAGAAcCTTCCATTTTGGGGGGGAGCTTACTTTTACTCTGTtTTTACAcATAGCAACCCACCTCTGAGAGGCCCCAGCCATAGACCCTACAGAGACAAGGCAGGCCTGACCCTAGCTATAATACACCCCTGAGACATGCTCTGAAACCGGGGGTGGGTGGGGACAGGGCAGGhCTGACCCCATTTGAAGGGGGCTCAGGCTCCCTAnCCAAAGATCTAGCCCTGTAGGGGTGTGTGTGTGTTTTCAAAGnGnCTTcACACTAGCACACTGGCTTGGAACTCGCTGTGTAGTCTAGAACTCATGACAAcTCTGTGCCCTGAGTTTCTCAAGGCTGGGATtACAGCCACCTGTAGCACCCGACCTTCTGTAGTGGAAAGCCCTTACCTTTTCCCCCAAGACTGGGAnCAGTcAACGCTTAGATTGTCAgGTGTtTTCAGTcAGTCTCCCTCCTCCCCCAGCTGCTCCTCAgCCAGgAG

Fig. 8B. (continued 3)

AgGAGGcAGGACcAcTcaAGGAGcAGGAACcAcCGgTgAgGTcCCTCCcAgAGcAGGcGGGcCCTGTcAAGTCTCCAc  
GACCCATTTATAcATTTCAGAcATCTCTACGGGTcACcATCGAAcGCCcAAGGcAAGAcAcAcAGTGTGGGTCTCTGG  
GGATTCGTCTGGcAGcCTGTGAACtTGGCCcAGAcAATAGGATGTCTCTCGCCCTAAAGTGATGGCTTCTGCCCCCTTCA  
CCTCTTTAGcAGAAtGGGTcAGAAcTAAAGTTAGAcTTCCcATTCCTCCcTCTGGcAAGAGGcAGcGTGTTCAATGTCTCCAGAG  
AAAcAcATTCCTCCtCnnCTTCTGGTTAAATCCAcCCCGTCTCCGTCTGAGGGcAACAACCCcATTTnAGTTGGCGGGT  
TCAcTGGTGTCTCCcCnCCAGGGCCAcAAATnAGnTTGTCTCAGGAATTATAcAnAcTAAAGTnAAAGGAnGT  
TCnnCCCTCCAAATTTTGGTTAAcCAAAcCAAGAnCnnTTGCCAGAGGGCTnTTAAAnAcAAATAAATTCaAGGAAC  
AACCCTCTTAGGCATTGnGTTGATCnCCGTGTCTGGcAcAcAcCATGGGTGGGTGGGTGCATGGTTAAAcAGAcAGCAAT  
GGATGTcATGTGTCTcAGAcGTATAcAcGTGAcAcCGTcAGAGAGGcAcGGnTCATGTCTTTcATAGAAcATGTGGATATT  
AGTGTGGAGAcAAATAGAAATTCATGAATTAGGGAATGTGTTCAGTGGCCCTGAGGAAGGGcAcAGTTTAGAAGAcAGCA  
TAGcAGAGAcCTTAGcATGGCTTCcAGAGGGATcAcAGAcCGCCCTGTCTTGGCTTCTTAAAGGcAcAGTTTAGAAGAcAGCA  
GCCTTAGAAGTTAcAGGGATCTcATAAGGGcATGGTGGAGGGCTGAGGGcAcAGCTTCcAcAGAGcCCTTCcAGGAcAGAc  
AGTCTTAGTTCCcAAcNtCTCTAcAGAACTGGAGGGTcAcCnTCAGAGcAGGAGGTAcAGTcAcCTTCGTGTcAGTCC  
TGCCAcAGGGcAGTGAACAcTCCAcCGTCTGGTCCcAcAGGTCTGTAAATGAAGGGGTGAAGTCTTTGTGTGAATACTA  
GAGCCTGGcAGTGGAGTCTCCcCTTAAATCCcAGCAATGGGGAGGTGAGGGcAGAcTCTGTGAGTTCAAGGTcAGCCAG  
GGCTAcAcAAAGAAATCCTGTCTCAAAcAAAcAAAGATGGATCC

# MAP OF THE HUMAN UCP2 GENE clone hUCP2-g2



\* 3,8 kb Bam HI fragment containing exon1(sequence 2), intron 1, exon 2 (sequence 3) and a part of intron 2.

\*\* the 5' region upstream the +1 site encompasses 3kb DNA or more.





**Fig. 10C.**

sequence 3

ATTGGCCTTGGGNAGGGCCNGGTTCCCGAGNCCTTTCCAAACTTTCTTNA  
CAGCCCCGACGNGGGGACNTAAGCCAATTTCCGGGGAGAGTTTNTGGTC  
CCANTGGNGACCCCCCTTNAAAAATTTTCCNATTNACCAAAGTNTGATGG  
ACTGNGTTAGGGGGTGCTTATATAGAGTACTGAGTGTAACAAAAGCAGA  
AGTCTGGATGAGAACCCAATTTGTGATATTAAGCAGGTGGGGTGGGGGT  
GGGGAGTGTACCTAGGTTCAATTTCCGCCNTGCTTNTCCCCCTTTCCAGT  
GTGTGCACTTAACCAGTCCCTGGGCCCTGTTCCCCATCCCCCTCCAAG  
GCATGGATTGGGTGGGCTTGTGTGTCTTGGGGCAGGTGGCCCTTTCTAA  
ACTCTCTGCCTTTGCTCACCCACAGGACACATAGTATGACCATTAGGTG  
TTTCGTCTCCACCCATTTTCTATGGAAAACCAAGGGGATCGGGCCATG  
ATAGCCACTGGCAGCTTTGAAGAANGGGACACCTTTAGAGAAGCTTGAT  
CTTGGAGGCCTCACCGTGAGACCTTACAAAGCCGGGTAAGAGTCCAGTC  
CAAGGAAGAGGTCTCTTGCTGCCTCCTAACCTGTGGTCTAGGGGCAGG  
AGTCAGCAGGGCATTAACAAAAATAATTACCATCCCCACCCCGCAGT  
GAAGTGGCTCTTTCCAGTTCACAGAGCACTCTCACACCTCCCCGCTCTC  
ATTCTGGCCCTTCAGCTGACTCGGACAAGCCAAGGATCTTGGTCCCCAT  
TTTATAAAGGAGAAAAGTGAAGGCCNCGTGTNACAGTGATTGGCCCCAA  
GTCATCCCGGGAGCCAGCAGAAGAGCTAGGACAGGAACCTATTGTTCTA  
ACTTCATATTGATGCTAGCTTTTGACTATCCCTGAAACCGAAATTGGTT  
ATCAGCCCGGCTCTGAAAGTGGTTATTTGCTGGGGACTGTTAAATAGGA  
TTAACTATTTCTATTCCTGCATTTTAATTGCTGTTAGTAGGGCAGCTTA  
CCACCCTCC

**Fig. 10D.**

sequence 4

TTCAGTGAAGCTCGTGGGCAAGCTGGAGGGAAACAGGAATAAGTTCAGGC  
CCTGTTcTATAGGTCCCAGTGTAGTTGCTATGGTGAGTATcTTCATTTCC  
CTGCTTGCCCCAGCCACcTGGAGTGAGAAGCCCAAGAGGAAGCTGGGTGA  
GCTGTTTGTTCATGGGTCTCTGTGTTTACAGCTGACTCCCTTCACCAG  
CCAGCCCTTTACCTGAGCCCCAGCAACAAAGGCAGTCAGGCGGGGCTCA  
AAGCAGCTGCTCCAATGAAGTCAAAGAAATAAGCTCAGGGGAAGAAGCAG  
GTCACCCTCCCCACTAGGGTGCTGGGCTCACTTCCTCCTGGGGCAGTGG  
AGGAGGGTGTGGTTCCAACCTCAGAACAAATGGGGCTTTTGGTTTACTTT  
ATCACTCTTCACAGCTCTGACCTGGACCCCTCATGCCCTGCCTGTCTGT  
GGTGTAAGTGCGGATCCCCCTAAGTTGGAGGAAAGGAACTGGCCCAAAC  
AAAAAGGAGAGCAGTTTTCTCTGCATCACATGGTAGGCCAGGAGGAGTCT  
AATGCCCCAGAGTTTACTCTCAGCCCCCAAATCACCTAGCTAAATGTTA  
CCTTATCTAAGAAGTCTTAGGTTTTTTGGGGTTTTTTTTTTTTTTTTT  
GAGACAAGGTCTCACTCTCTCACCAGACTGGAGCACAGTGGCACAATCA  
CAGCTCACTGCAGCCTCAACCTCCTGGGCTCAAGCAATCGTCCCAAGTAG  
CTGGGACTATAGGCCTGCACCACCATGTCCAGCTAATTTATTTTATTTA  
TATTTTTTAgACAGGGTCTCATTATGTTGCCCTGGCTGGTCTTGAACCTC  
TGGGTTCAAGcCAGTCTCCACCTCTGCCTCCCAAAGTGCTAGGTTTTTT  
TtGTTTGTGTTTGTGTTTGTGTTTGTGTTTGTGTTTGTGTTTGTGTTT  
TAGGCTGGAGTGAGTGGCACgATCTCAGcTACTGCAACcTCCACcTCcT  
GGGTTCAAGTGATTcTCTGCCTCAGCTCCTAAGTAgtTGGGAATACAG  
GCGTGTGCCAACACACCCAGCTCATTTTTtGTATTTTTTAgCGGAgATGGGG


**Fig. 10D.** (continued 1)

TTTTGCCATGTTGGCCAAGCTGGTCTCAAACCTCCTGA<sub>c</sub>CTCAGGTGATTC  
GCCCCGCTCAGCCTCCCAAAGTGCTGGGTTTACAGGCGTGAGCCACCACA  
CCCAGCCCAAGAAgT<sub>c</sub>TTTT<sub>c</sub>TGAT<sub>c</sub>ACCCAcTCTT<sub>c</sub>CTCTCCCAAt  
GGCATtAgTtGTTCCCTCCTTTGCATTTTGAGAGTATGTCctGTAAGCCC  
cAAATG<sub>c</sub>AgCtTGAATCATCTGCCCATC<sub>c</sub>ACCCCCCTGTGCCCAACAGTAA  
GCCTCCTCTAGAGTAGATACTAtCTCCTGCATCTCAGTGAACCACTGCC<sub>c</sub>  
AgCAAAGCAGTCTTGCTAAAACAA<sub>t</sub>GACT<sub>c</sub>TAGAg<sub>a</sub>TCCTAAGCTGTGtG  
AGAGCTGGAGGAGAGAATTAGAcTGATG<sub>t</sub>CTGGGAAGGGATTGAATTAG  
TCATCTTGtACCTTTTCTTCTTGACTTAAGTTCCAGACCTGTAGCAACCA  
TTCCTGCTTAGACATCCAGAACATAAGCCTAtGGgt<sub>c</sub>tGTGCCTGTTGGG  
TCTTAGTCTGGGTGAAACTTTTCTCTACTTCTGTGAGCtCTCCAGATGAA  
CCACAGAAGCAGGAATGTGGGCATCATCAGTGAAATCTCTGCATACAGCA  
GACAAAGGGCTGGTCCAGTGGCTGTTTATGAGGCAGCGCTAGGAGAGCTC  
TGATCCAGACTCTCCCTGCAGTGAAAGGGAGGGAGCCCTTCATGAAGTAT  
TGACTGCTTGAGCAGGAATTGCTTCACCAGCACCTAACTGAGTGCCTCTC  
GAGCTCACATCGGTTTTCCCTCATGAGGCCACTTGGAGTCTTGCTGAGGG  
ACTTGgTTCTATTAGGGAAGGTGAGTTTGGGGATGGTGAGCAGGGAGGGC  
CTGGGGACATTGTGGCTAATGGGGCTTTTCTCCTCTTGgCTTAGATTCCG  
GCAGAGTTCCCTCTATCTCGTCTTGTTGCTGATTAAAGGTGCCCTGTCTC  
CAGTTTTTCTCCATCTCCTGGGACGTAGCAGGAAATCAGCATCATGGTTG  
GGTTCAAGGCCACAGATGTGCCCCCTACTGCCACTGTGAAGTTTCTTGGG  
GCTGGCACAGCTGCCTGCATCGCAGATCTCATCACCTTTCCTCTGGATAC  
TGCTAAAGTCCGGTTACAGGTGAGGGGATGAAGCCTGGGAGTCTTGATGG  
TGTCTACTCTGTTCCCTCCCCAAAGACACAGACCCCTCAAGGGCCAGTGT  
TTGGAGCATCGAGATGACTGGAGGTGGGAAGGGCAACATGCTTATCCCTG  
TAGCTACCCTGTCTTGGCCTTGcAGATCCAAGGAGAAAGTCAGGGGGCCAG  
TGCGCGCTACAGCCAGCGCCCAAGTACCGCGGTGTGATGGGCACCATTCTG  
ACCATGGTGCCTACTGAGGGCCCCCGAAGCCTCTACAATGGGCTGGTTGC  
CGGCCTGCAGCGCCAAATGAGCTTTGCCTCTGTCCGCATCGGCCTGTATG  
ATtCTGTCAA<sub>a</sub>CAGTtCTACACCAAgGGCTCTGAgC<sub>g</sub>TGAgTATGgAgCA  
aGGGTGTAGGCCCTTtGGCCCTTTTt<sub>t</sub>CTCagTGatGATtGATCTtAGTt  
CATtCAgCCATATAgTtTtTtAGgCCC<sub>c</sub>AcgaT<sub>c</sub>CcTAGgAAGatCAGGG  
GgAACAGAGAACTGGAAGGGGGCCCTGGTCTCCACATAGTTCCTAAGCAC  
CTGGGCTATACCAGG<sub>c</sub>TCTGAGCAGGgCGTCATCCCATCACAGTCTTCAA  
CACCACCTTGGGAGTAGGTAGTATCATCCAGTGTATAGAaGAAgAgAC  
TGAGGTGGGAAGGCAgTGGGTAgAGTGGGGACTTGGCCAGGGGCACACAG  
TAGAGAGCCAGAAAACACACAGTAGAGAGCCAGGACACTCGTCTCTAAGG  
CCAGCGTTCTTCCCTTTTCAcCTCCTtAGtAtGCCAtGCCAACC<sub>c</sub>TCCATT  
TTACACATGACGAAACaGaGCCCCAAACAAAAGGTTGTCTTTCCAGATC  
ACATGGCAGGAAGAAGTAAAGCTGAcCTGAGATCCCAAGTCTTAGGAATC  
CCAGTCTCAGAAAGCCACTTCTCTCTGAGCCTTGGTTTTTACATTGTGTC  
AGATGGAaATGATTGTGATTTCTCAGGGCTGTTGAGCAGGTAAATGAAAA  
TGTTTTATGAAAGAAAGCACCAAGTTTCATTTTGGTCTTAGCCCTTGCTA  
TGTCCCTAGCAAGAAGTAGATATTCATAGGGATATTTTGTtTGATGTGAG  
GAGTTCTTACAGCAAGAGCTTGTAGAAGGCCAAAAGCTTCTGGATTCTAA  
TCCCAAAAGCAGGAGATGACAGTGACAGGGTGGTTTTTGGTGAGGAGAGAT  
GAGGTAGAAAATGAGTGCAAGCCCGCTGGCCACTGACCCCATGGCTCGCC  
CACAGATGCCAGCATTGGGAGCCGCTCCTAGCAGGCAGCACACAGGTG  
CCCTGGCTGTGGCTGTGGCCCAGCCACGGATGTGGTAAAGGTCCGATTC

**Fig. 10D. (continued 2)**

CAAGCTCAGGCCCCGGGCTGGAGGTGGTCGGAGATACCAAAGCACCGTCAA  
TGCTTACAAGACCATTGCCCGAgAGGAAGGGTTCCGGGGCCTCTGGAAAG  
GTGTGTACCAGTTGTTTTCCCTTCCCTTTTCTCCTCCCCGATACTCTG  
GTCTCACCCAGGATCTTCTCCTCCTACAGGGACCTCTCCCAATGTTGCT  
CGTAATGCCATTGTCAACTGTGCTGAGcTGGTGACCTATGAcCTCATCAA  
GGATGCCCTCCTGAAAGCCAACCTCATGACAGGTGAGTCATGAgGTAgAc  
gGTGCTgGGTCTCACCTTCCCCcATGcCaggAgCAGgTGCGGGGgTcTA  
gCTGAcACcAgAagAccacATCTTTTCATCCTATTTGCCCTTTGCAGGGA  
GAGTAaGaAtATCTCTTACTTGCCATATtGAAGCCAATTGGGATGAAgCTC  
CCACTTTGCACaTtGAGgAAcTGAgGCTAgAtTGGCAAAATGAcTCTTtC  
AGgTCCTCAGAAGATGTCTCAGCTGGAGTCCCTGtCtgTTTTTgTTTTT  
TgtttGtttGttTTTTGttTTTTtGaGAtAGaGTcTCAcTcTGttACCC  
GtGTAATcTCAGCTCACTGCAACCTTCTCCTccTGGGTTCAGcGATTCT  
TGTGCCTCAGCCTCCCgAgTAGcTGGGATGACAGGTGTGCACCgCACAC  
tGGCTAATTTTTGTATTTTTAGTAGAgAtGGAGTTTCACCATGTTAGCCA  
GGCTGGTCTcGAAcTCTGGCCTCaAGTGATCTGCCCACCTTGGCCTCCC  
AATGTGCTGGGATTACAGGTGTgAGCCTcTGCGCCCCAtCcTcTTGtTtG  
tTtTTTGAgACAGGGTCTtGcTCGGTtGCCAgGctGGAgTGCAGTGGGG  
tgATTAATGGcTCATtGcAgCctcgACCTCCcTGAcTCAAgCAATCcTCC  
CACcTCaGCCTCcTGAgTAGcTGGGGCTGActAcaGGCaTgCACACTGTg  
CCTGGCTaATTTTTGTaTTTTGTAgAgACAGGGTTTTTGCCATGTTACCC  
AgTctGGTCTTgAAcTcTGGGCTCaaGtGATCcaCCAcCtcggCCTcc  
AAAAGAAGTCCTGGATTACAGGCATGAGACATTGTGCCCAGCCTCTCTGT  
CTCTTTAAATCATGAAACTCGTAGCTACTTAAGTAATTCTCCTGCCTT  
CTGGAATGATGGGTGAAGATCTTGACTGCCTTGCCCTGCTCCTCCTTGGCA  
GATGACCTCCCTTGCCACTTCACTTCTGCCTTTGGGGCAGGCTTCTGCAC  
CACTGTcATCGCcTCCCCTGTAGACGTGGTCAAGACGAGATACATGAAcT  
CTGCCCTGGGCCAGTACAGTAGCGCTGGCCACTGTGCCCTTACCATGCTC  
CAGAAGGAGGGgCCCCGAgCCTTctACaAAGGgTGAGcCTcTGgTCcTCC  
CcACCCAGTTCAGGCcTcTTGGcTATGCATGTCTATTATGGGTGGGAGAG  
AACCACCTGGAAGTGAGTAGCAGCCAAGTGTGACTATTTCTGATCCTGGT  
CGTGGCATTTCACCAGCATTCAcCTATCcCcTtAaTtCcTtCcTCCcagA  
aTtGcTAcCATCAcTGTtATTAGGTGTtAAATGGAGACTCaAaGGGAaT  
tCATGCTTATAGCCaAGCAGcTGTGAGCTCAGtTCAttGAgTCcTCCAG  
CCTCCTTtgGGAcAGAgCAAcTGGGTtGGAtTGAATAcCAGGCCcAGTGA  
GGGAAGTGGGAGGTGGAGGTGCCCCCATGACCTGTGATTTTTCTCCTCTA  
GGTTCATGCCCTCCTTTCTCCGCTTGGGTTCCTGGAACGTGGTGATGTTc  
GTCACCTATGAGCAGCTGAAACGAGCCCTCATGGCTGCCTGCACCTCCCG  
AGAGGCTCCCTTCTGAGCCTCTCCTGCTGCTGACCTGATCACCTCTGGCT  
TTGTCTCTAGCCGGGCCATGCTTTCTTTCTTCTCCTTCTTCTCTCCCT  
CCTTCCCTTCTCTCCTTCCCTCTTTCCCCACCTCTTCTTCCGCTCCTTT  
ACCTACCACCTTCCCTCTTTCTACATTCTCATCTACTCATTGTCTCAGTG  
CTGGTGGAGTTGACATTtGACAGTGTGGGAGGCCTCGTACCAGCCAGGAT  
CCCAAGCGTCCCGTCCCTTGGAAAGTTCAGCCAGAATcTTcGTCTTCCCC  
CCGACAGCCcAGCCTAGCCCAcTTGTcATCCATAAAGCAAGCTCAACCTT  
GGCGTCTCCTCCcTCTCTTGTAGCTCTTACCAGAGGTCTTGGTCCAATGG  
CCTTTTTTGGTACCTGGTGGGcAGGGGAGGAACCACCTGACTTTGAAAATG  
GGTGTGATCCACCTTCCACCTCCAGCATCCAATCTGAAGCCCGTGTAGGT



*Fig. 11.*

# INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>16</u> , lines <u>21-25</u>	
<b>B. IDENTIFICATION OF DEPOSIT</b> <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depositary institution <p style="text-align: center;">American Type Culture Collection</p>	
Address of depositary institution (including postal code and country) <p style="text-align: center;">12301 Parklawn Drive Rockville, MD 20852, USA</p>	
Date of deposit <p style="text-align: center;">April 18, 1997</p>	Accession Number <p style="text-align: center;">97993</p>
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indications are not for all designated States)	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

<p style="text-align: center;">For receiving Office use only</p> <div style="border: 1px solid black; padding: 5px; margin-bottom: 5px;"> <input type="checkbox"/> This sheet was received with the international application       </div> <div style="border: 1px solid black; padding: 5px; min-height: 40px;">         Authorized officer       </div>	<p style="text-align: center;">For International Bureau use only</p> <div style="border: 1px solid black; padding: 5px; margin-bottom: 5px;"> <input checked="" type="checkbox"/> This sheet was received by the International Bureau on:  <p style="text-align: center;">23 February 1998 (23.02.98)</p> </div> <div style="border: 1px solid black; padding: 5px; min-height: 40px;">         Authorized officer  <p style="text-align: center;">Ting Zhao</p> </div>
---	---

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US97/06864

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : 435/69.1, 252.2, 254.11, 320.1, 325; 514/44; 530/350; 536/23.5

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/69.1, 252.2, 254.11, 320.1, 325; 514/44; 530/350; 536/23.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	FLEURY et al. Uncoupling protein-2: a novel gene linked to obesity and hyperinsulinemia. Nature Genetics. 15 March 1997. Vol. 15, No. 3, pages 269-272, especially the Abstract and Figure 1.	9, 10, 12, 13
X	RAIMBAULT et al. Mus musculus UCP2 mRNA, complete cds. GenBank, Accession Number U69135. 30 October 1996. See the entire document, especially residues 813-827.	9, 12
X	HILLIER et al. yf39c01.r1 Homo sapiens cDNA clone 129216 5' similar to SP:UCP_RABIT P14271	12, 13, 15, 16
Y	MITOCHONDRIAL BROWN FAT UNCOUPLING PROTEIN. EST-STS, Accession Number R11086. 11 April 1995. See the entire document, especially under Source.	1, 2, 4, 177

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

•	Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A"	document defining the general state of the art which is not considered to be of particular relevance		
"E"	earlier document published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means		
"P"	document published prior to the international filing date but later than the priority date claimed	"Z"	document member of the same patent family

Date of the actual completion of the international search

03 AUGUST 1997

Date of mailing of the international search report

03 SEP 1997

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

BRIAN LATHROP

Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT.

International application No.  
PCT/US97/06864

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	CHARNOCK-JONES et al. Extension of incomplete cDNAs (ESTs) by biotin/streptavidin-mediated walking using the polymerase chain reaction. Journal of Biotechnology. 30 June 1994. Vol. 35, pages 205-215, see the entire document.	1, 2, 4, 15-177
Y	US 5,453,270 A (BILLS) 26 September 1995, column 2, line 24, through column 3, line 14, and column 3, line 53, through column 4, line 19.	1, 2, 4, 15-17
Y	KOZAK et al. The mitochondrial uncoupling protein gene. The Journal of Biological Chemistry. 05 September 1988. Vol. 263, No. 25, pages 12274-12277. See the entire article.	1, 2, 4, 15-17



# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US97/06864

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☒ Claims Nos.: 11 and 14  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  
  
Please See Extra Sheet.
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  
1-4 and 9-10,12-13,15-17

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US97/06864

## A. CLASSIFICATION OF SUBJECT MATTER: IPC (6):

A61K 48/00; C07K 14/47; C12N 1/15, 1/21, 5/10, 15/09, 15/12; C12P 21/02

## B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

GenBank, EST-STS, issued US patents, USPAT, MEDLINE, CAPLUS, WPIDS  
search terms: GenBank Accession Numbers U76367 (H. sapiens UCP2) and U69135 (M. musculus UCP2), ucp2, uncoupling protein, mitochondria, glucose, hyperinsulinemia

## BOX I. OBSERVATIONS WHERE CLAIMS WERE FOUND UNSEARCHABLE

2. Where no meaningful search could be carried out, specifically:

Claims 11 and 14 are unsearchable to the extent that they require reference to the specified sequences from the Figures. Because applicant has not furnished a computer-readable sequence listing, no meaningful search of the sequences per se can be carried out by this Authority. However, the subject matter of the claims has been searched to the extent possible with reference to the balance of the description.